

TLR11 Antibody

Catalog # ASC10236

Product Information

Application	WB, IF, ICC, E
Primary Accession	Q6R5P0
Other Accession	NP_991388 , 45429992
Reactivity	Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	105873
Concentration (mg/ml)	1 mg/mL
Conjugate	Unconjugated
Application Notes	TLR11 antibody polyclonal antibody can be used for detection of TLR11 by Western blot at 0.5 to 2 μ g/mL. Antibody can also be used for immunocytochemistry starting at 10 μ g/mL. For immunofluorescence start at 10 μ g/mL.

Additional Information

Gene ID	239081
Other Names	TLR11 Antibody: Gm287, Gm287, Toll-like receptor 11, toll-like receptor 11
Target/Specificity	Tlr11;
Reconstitution & Storage	TLR11 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.
Precautions	TLR11 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	Tlr11 {ECO:0000303 PubMed:14993594, ECO:0000312 MGI:MGI:3045226}
Function	Participates in the innate immune response to microbial agents. Acts via MYD88 and TRAF6, leading to NF-kappa-B activation, cytokine secretion and the inflammatory response.
Cellular Location	Membrane; Single-pass type I membrane protein


Background

TLR11 Antibody: Toll-like receptors (TLRs) are signaling molecules that recognize different microbial products during infection and serve as an important link between the innate and adaptive immune responses. These proteins act through adaptor molecules such as MyD88 and TIRAP to activate various kinases and transcription factors. TLR11 is one of three mouse TLRs that lack a human ortholog. It is activated specifically by uropathogenic bacteria, and mice lacking TLR11 showed a much greater susceptibility to uropathogenic infections, indicating a potentially important role for TLR11 in preventing infections in the urogenital system.


References

Vogel SN, Fitzgerald KA, and Fenton MJ. TLRs: differential adapter utilization by toll-like receptors mediates TLR-specific patterns of gene expression. *Mol. Interv.* 2003; 3:466-77.
Takeda K, Kaisho T, and Akira S. Toll-like receptors. *Annu. Rev. Immunol.* 2003; 21:335-76.
Janeway CA Jr. and Medzhitov R. Innate immune recognition. *Annu. Rev. Immunol.* 2002; 20:197-216.
O'Neill LAJ, Fitzgerald FA, and Bowie AG. The Toll-IL-1 receptor adaptor family grows to five members. *Trends in Imm.* 2003; 24:286-9.


Images



Western blot analysis of TLR11 in RAW264.7 cell lysates with TLR11 antibody at (A) 0.5, (B) 1, and (C) 2 µg/mL.



Immunocytochemistry of TLR11 in RAW264.7 cells with TLR11 antibody at 10 µg/mL



Immunofluorescence of TLR11 in Raw264 cells with TLR11 antibody at 10 µg/mL.