

West Nile Virus Envelope Antibody

Catalog # ASC10289

Product Information

Application	E
Primary Accession	P06935
Other Accession	NP_776014 , 27735302
Reactivity	Virus
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	380110
Concentration (mg/ml)	1 mg/mL
Conjugate	Unconjugated
Application Notes	West Nile virus envelope antibody can be used for the detection of the West Nile virus envelope protein in ELISA. It will detect 10 ng of free peptide at 1 μ g/mL.

Additional Information

Gene ID	912267
Other Names	West Nile Virus Envelope Antibody: Genome polyprotein, Core protein, NS1, Envelope protein
Target/Specificity	WNVgp1;
Reconstitution & Storage	West Nile Virus Envelope antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.
Precautions	West Nile Virus Envelope Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	POLG
Function	[Capsid protein C]: Capsid protein C self-assembles to form an icosahedral capsid about 30 nm in diameter. The capsid encapsulates the genomic RNA (Probable). Plays a role in virus budding by binding to the cell membrane and gathering the viral RNA into a nucleocapsid that forms the core of a mature virus particle (By similarity). During virus entry, may induce genome penetration into the host cytoplasm after hemifusion induced by the surface proteins (By similarity). Can migrate to the cell nucleus where it modulates host functions (By similarity). Overcomes the anti-viral effects of host EXOC1 by sequestering and degrading the latter through the proteasome

degradation pathway (PubMed:[23522008](#)).

Cellular Location

[Capsid protein C]: Virion {ECO:0000250|UniProtKB:P17763}. Host nucleus {ECO:0000250|UniProtKB:P17763}. Host cytoplasm. Host cytoplasm, host perinuclear region. Host nucleus, host nucleolus {ECO:0000250|UniProtKB:Q01299} [Small envelope protein M]: Virion membrane {ECO:0000250|UniProtKB:P03314}; Multi-pass membrane protein {ECO:0000250|UniProtKB:P03314}. Host endoplasmic reticulum membrane {ECO:0000250|UniProtKB:P03314}; Multi-pass membrane protein. Note=ER membrane retention is mediated by the transmembrane domains. {ECO:0000250|UniProtKB:P03314} [Non-structural protein 1]: Secreted {ECO:0000250|UniProtKB:P17763}. Host endoplasmic reticulum membrane; Peripheral membrane protein; Luminal side {ECO:0000250|UniProtKB:P17763}. Note=Located in RE-derived vesicles hosting the replication complex. {ECO:0000250|UniProtKB:Q9Q6P4} [Serine protease subunit NS2B]: Host endoplasmic reticulum membrane; Multi-pass membrane protein {ECO:0000250|UniProtKB:P17763} [Non-structural protein 4A]: Host endoplasmic reticulum membrane {ECO:0000250|UniProtKB:P14335}; Multi-pass membrane protein {ECO:0000250|UniProtKB:P17763}. Note=Located in RE-associated vesicles hosting the replication complex {ECO:0000250|UniProtKB:P17763} [RNA-directed RNA polymerase/Methyltransferase NS5]: Host endoplasmic reticulum membrane; Peripheral membrane protein; Cytoplasmic side. Host nucleus. Host cytoplasm {ECO:0000250|UniProtKB:P14335}. Note=Located in RE-associated vesicles hosting the replication complex. NS5 protein is mainly localized in the nucleus rather than in ER vesicles (By similarity) Shuttles between the cytoplasm and nucleus (By similarity) {ECO:0000250|UniProtKB:P14335, ECO:0000250|UniProtKB:P17763}

Background

West Nile Virus Envelope Antibody: West Nile Virus (WNV) is a member of the Flaviviridae, a plus-stranded virus family that includes St. Louis encephalitis virus, yellow fever virus, and Dengue virus. WNV was initially isolated in 1937 in the West Nile region of Uganda and has become prevalent in Africa, Asia, and Europe. It has rapidly spread across the United States with cases being observed in every continental state. Virus particles consist of a dense core made up of the core/capsid protein encapsulating the RNA genome surrounded by a membrane envelope embedded with envelope and matrix proteins. While the viral core protein is thought to contribute to the WNV-associated inflammation via apoptosis induced through the caspase-9 pathway, the highly glycosylated envelope protein plays a major role for WNV entry into target cells as this entry can be inhibited by using a recombinant domain III from the envelope glycoprotein. The WNV receptor has recently been identified as $\alpha v \beta 3$ integrin.

References

- Gould LH and Fikrig E. West Nile virus: a growing concern. *J. Clin. Invest.* 2004; 113:1102-7.
- Yang JS, Ramanathan MP, Muthumani K, et al. Induction of inflammation by West Nile Virus capsid through the caspase-9 apoptotic pathway. *Emerg. Infect. Dis.* 2002; 8:1379-84.
- Chu JJ, Rajamanonmani R, Li J, et al. Inhibition of West Nile virus entry by using a recombinant domain III from the envelope glycoprotein. *J. Gen. Virol.* 2005; 86:405-12.
- Chu JJ and Ng ML. Interaction of West Nile virus with $\alpha v \beta 3$ integrin mediates virus entry into cells. *J. Biol. Chem.* 2004; 279:54533-41.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.