

# **BAP29 Antibody**

Catalog # ASC10383

## **Product Information**

Application WB, E
Primary Accession Q9UHQ4

Other Accession NP\_001008405, 56549093
Reactivity Human, Mouse, Rat

Host Rabbit
Clonality Polyclonal
Isotype IgG
Calculated MW 28320
Concentration (mg/ml) 1 mg/mL
Conjugate Unconjugated

Application Notes Bap29 antibody can be used for the detection of Bap29 by Western blot at 1 -

2 [g/mL.

### **Additional Information**

**Gene ID** 55973

Other Names BAP29 Antibody: BAP29, BAP29, B-cell receptor-associated protein 29,

BCR-associated protein 29, B-cell receptor-associated protein 29

Target/Specificity BCAP29;

**Reconstitution & Storage** BAP29 antibody can be stored at 4°C for three months and -20°C, stable for

up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high

temperatures.

**Precautions**BAP29 Antibody is for research use only and not for use in diagnostic or

therapeutic procedures.

#### **Protein Information**

Name BCAP29

Synonyms BAP29

**Function** May play a role in anterograde transport of membrane proteins from the

endoplasmic reticulum to the Golgi. May be involved in CASP8- mediated

apoptosis (By similarity).

**Cellular Location** Endoplasmic reticulum membrane; Multi-pass membrane protein

## **Background**

BAP29 Antibody: Bap29 and the related protein Bap31 are endoplasmic reticulum (ER) and ER-vesicle membrane proteins and members of the B-cell receptor-associated protein family. These two proteins are highly homologous and can form homo- and heterodimers. Both Bap29 and Bap31 interact with membrane-bound immunoglobulins (mIgs), such as IgM and IgD, which with Ig-alpha/Ig-beta heterodimers form B cell antigen receptors. Binding of the Bap29/Bap31 heterodimer correlates with the ER retention of non-Ig-alpha/Ig-beta bound mIg complexes, suggesting that Bap29 and Bap31 may act as chaperones transmembrane regions of various proteins. Bap29 possesses multiple isoforms.


#### References

Kim KM, Adachi T, Nielsen PJ, et al. Two new proteins preferentially associated with membrane immunoglobulin D. EMBO J.1994; 13:3793-800.

Ng F, Nguyen M, Kwan T, et al. p28 BAP31, a Bcl-2/Bcl-XL- and procaspase-8-associated protein the endoplasmic reticulum. J. Cell Biol.1997; 139:327-38.

Schamel WW, Kuppig S, Becker B, et al. A high-molecular-weight complex of membrane proteins BAP29/BAP31 is involved in the retention of membrane-bound IgD in the endoplasmic reticulum. Proc. Natl. Acad. Sci. USA2003; 100:9861-6.

## **Images**



Western blot analysis of Bap29 in human heart tissue lysate with Bap29 antibody at (A) 1 and (B) 2 µg/mL.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.