

TLR9 Antibody

Catalog # ASC10393

Product Information

Application	WB, IF, E, IHC-P
Primary Accession	Q9NR96
Other Accession	AAH32713 , 21595773
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	115860
Concentration (mg/ml)	1 mg/mL
Conjugate	Unconjugated
Application Notes	TLR9 antibody can be used for detection of TLR9 by Western blot at 0.5 to 1 μ g/mL. Antibody can also be used for immunohistochemistry starting at 2 μ g/mL. For immunofluorescence start at 10 μ g/mL.

Additional Information

Gene ID	54106
Other Names	TLR9 Antibody: CD289, Toll-like receptor 9, toll-like receptor 9
Target/Specificity	TLR9;
Reconstitution & Storage	TLR9 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.
Precautions	TLR9 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	TLR9
Function	Key component of innate and adaptive immunity. TLRs (Toll- like receptors) control host immune response against pathogens through recognition of molecular patterns specific to microorganisms. TLR9 is a nucleotide-sensing TLR which is activated by unmethylated cytidine- phosphate-guanosine (CpG) dinucleotides (PubMed: 14716310). Acts via MYD88 and TRAF6, leading to NF-kappa-B activation, cytokine secretion and the inflammatory response (PubMed: 11564765 , PubMed: 17932028). Controls lymphocyte response to Helicobacter infection (By similarity). Upon CpG stimulation, induces B-cell proliferation, activation, survival and antibody production (PubMed: 23857366).

Cellular Location	Endoplasmic reticulum membrane; Single-pass type I membrane protein {ECO:0000250 UniProtKB:Q9EQU3}. Early endosome membrane. Lysosome {ECO:0000250 UniProtKB:Q9EQU3} Cytoplasmic vesicle, phagosome {ECO:0000250 UniProtKB:Q9EQU3}. Golgi apparatus membrane. Note=Relocalizes from endoplasmic reticulum to endosome and lysosome upon stimulation with agonist. Exit from the ER requires UNC93B1. Endolysosomal localization is required for proteolytic cleavage and subsequent activation Intracellular localization of the active receptor may prevent from responding to self nucleic acid. {ECO:0000250 UniProtKB:Q9EQU3, ECO:0000269 PubMed:14716310, ECO:0000269 PubMed:38169466}
Tissue Location	Highly expressed in spleen, lymph node, tonsil and peripheral blood leukocytes, especially in plasmacytoid pre-dendritic cells. Levels are much lower in monocytes and CD11c+ immature dendritic cells. Also detected in lung and liver


Background

TLR9 Antibody: Toll-like receptors (TLRs) are evolutionarily conserved pattern-recognition molecules resembling the toll proteins that mediate antimicrobial responses in *Drosophila*. These proteins recognize different microbial products during infection and serve as an important link between the innate and adaptive immune responses. TLR9 forms a subfamily along with TLR7 and TLR8 that recognize viral RNA and CpG DNA sequences and are localized in intracellular acidic compartments such as the phagolysosome. Unlike other TLRs which act through adaptor molecules such as TOLLIP, TIRAP, TRIF, and MyD88 to activate various kinases and transcription factors to respond to potential infection, TLR9 is strictly dependent on MyD88.

References



Takeda K, Kaisho T, and Akira S. Toll-like receptors. *Annu. Rev. Immunol.* 2003; 21:335-76.
Janeway CA Jr. and Medzhitov R. Innate immune recognition. *Annu. Rev. Immunol.* 2002; 20:197-216.
Wagner H. The immunobiology of the TLR9 subfamily. *Trends Immunol.* 2004; 381-6
Nishiya T and DeFranco AL. Ligand-regulated chimeric receptor approach reveals distinctive subcellular localization and signaling properties of the Toll-like receptors. *J. Biol. Chem.* 2004; 279:19008-17.

Images



Western blot analysis of TLR9 in mouse spleen cell lysate with TLR9 antibody at (A) 0.5 and (B) 1 µg/mL.

Immunohistochemistry of TLR9 in mouse spleen cells with TLR9 antibody at 2 µg/mL.



Immunofluorescence of TLR9 in mouse spleen cells with TLR9 antibody at 10 $\mu\text{g/mL}$.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.