10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

IL-32 Antibody

Catalog # ASC10399

Product Information

Application	WB, IF, E, IHC-P
Primary Accession	P24001
Other Accession	AAH09401, 14424787
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	26676
Concentration (mg/ml)	1 mg/mL
Conjugate	Unconjugated
Application Notes	IL-32 antibody can be used for the detection of IL-32 by Western blot at 5 - 10 g/mL. Antibody can also be used for immunohistochemistry starting at 10 g/mL. For immunofluorescence start at 20 g/mL.

Additional Information

Gene ID	9235
Other Names	IL-32 Antibody: NK4, TAIF, TAIFa, TAIFb, TAIFc, TAIFd, IL-32beta, IL-32alpha, IL-32delta, IL-32gamma, NK4, Interleukin-32, Natural killer cells protein 4, IL-32, interleukin 32
Target/Specificity	IL32; This antibody detects all isoforms of IL-32.
Reconstitution & Storage	IL-32 antibody can be stored at 4℃ for three months and -20℃, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.
Precautions	IL-32 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	IL32
Synonyms	NK4, TAIF
Function	Cytokine that may play a role in innate and adaptive immune responses. It induces various cytokines such as TNFA/TNF and IL8. It activates typical cytokine signal pathways of NF-kappa-B and p38 MAPK.
Cellular Location	Secreted.

Tissue Location	Selectively expressed in lymphocytes. Expression is more prominent in immune cells than in non-immune cells


Background

IL-32 Antibody: Interleukin-32 (IL-32) was initially identified as a transcript (NK4) that is selectively expressed in lymphocytes and NK cells and whose expression is increased following activation by IL-2. It was later re-isolated from an IL-18-treated lung carcinoma cell line and re-named IL-32. IL-32 is unusual in that it does not share sequence homology with known cytokine families and is highly expressed in immune tissues, existing in at least four differentially spliced isoforms. Because treatment of human monocytic and mouse macrophage cells with IL-32 induces several proinflammatory cytokines such as TNF-α, IL-8 and MIP-2, and because it is also induced in human peripheral lymphocyte cells after mitogen stimulation and in epithelial cells by IFNγ, it has been suggested that IL-32 may play a role in autoimmune and inflammatory diseases such as rheumatoid arthritis.


References

Dahl CA, Schall RP, He HL, et al. Identification of a novel gene expressed in activated natural killer cells and T cells. J. Immunol. 1992; 148:597-603.
Kim S-H, Han S-Y, Azam T, et al. Interleukin-32: a cytokine and inducer of TNF-a. Immunity 2005; 22:131-42.
Cagnard N, Letourneur F, Essabbani A, et al. Interleukin-32, CCL2, PF4F1 and GFD10 are the only cytokine/chemokine genes differentially expressed by in vitro cultured rheumatoid and osteoarthritis fibroblast-like synoviocytes. Eur. Cyto. Network 2005; 16:289-92.

Images




Western blot analysis of IL-32 in human spleen lysate with IL-32 antibody at (A) 5 and (B) 10 µg/mL shows two isoforms of IL-32.



Immunohistochemistry of IL-32 in human spleen tissue with IL-32 antibody at 10 µg/mL.

Immunofluorescence of IL-32 in Human Spleen cells with IL-32 antibody at 20 µg/mL.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.