

# TIM-1 Antibody

Catalog # ASC10421

### **Product Information**

Application	WB, IF, E, IHC-P
Primary Accession	<u>096D42</u>
Other Accession	<u>NP_036338</u> , <u>153085427</u>
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	39250
Concentration (mg/ml)	1 mg/mL
Conjugate	Unconjugated
Application Notes	TIM-1 antibody can be used for the detection of TIM-1 by Western blot at 1 - 2 [g/mL. Antibody can also be used for immunohistochemistry starting at 10 [g/mL. For immunofluorescence start at 20 [g/mL.

#### **Additional Information**

Gene ID Other Names	26762 TIM-1 Antibody: TIM, KIM1, TIM1, HAVCR, KIM-1, TIM-1, TIMD1, TIMD-1, HAVCR-1, Hepatitis A virus cellular receptor 1, Kidney injury molecule 1, HAVcr-1, hepatitis A virus cellular receptor 1
Target/Specificity	HAVCR1;
Reconstitution & Storage	TIM-1 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.
Precautions	TIM-1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

#### **Protein Information**

Name	HAVCR1
Synonyms	KIM1, TIM1, TIMD1
Function	Phosphatidylserine receptor that plays an important functional role in regulatory B-cells homeostasis including generation, expansion and suppressor functions (By similarity). As P- selectin/SELPLG ligand, plays a specialized role in activated but not naive T-cell trafficking during inflammatory responses (PubMed: <u>24703780</u> ). Controls thereby T-cell accumulation in the inflamed central nervous system (CNS) and the induction

	of autoimmune disease (PubMed: <u>24703780</u> ). Also regulates expression of various anti- inflammatory cytokines and co-inhibitory ligands including IL10 (By similarity). Acts as a regulator of T-cell proliferation (By similarity). May play a role in kidney injury and repair (PubMed: <u>17471468</u> ).
Cellular Location	Cell membrane; Single-pass type I membrane protein
Tissue Location	Widely expressed, with highest levels in kidney and testis. Expressed by activated CD4+ T-cells during the development of helper T-cells responses.

# Background

TIM-1 Antibody: The human form of TIM-1 was initially discovered as a membrane glycoprotein through which the hepatitis A virus can gain entry into a cell. It was also identified as kidney injury molecule 1 (Kim-1), a predicted adhesion molecule that is upregulated on the surfaces of kidney epithelia. It is also expressed on T helper 2 (Th2) cells of the immune system, and following the binding of its natural ligand TIM-4, stimulates T cell expansion and cytokine production in response to viral challenge. It has been suggested that hyperactivation of TIM-1 leads to an increased level of Th2 responsiveness and asthma susceptibility, and antibodies to TIM-1 may therefore be a novel approach to treating asthma.

## References

Feigelstock D, Thompson P, Mattoo P, et al. The human homolog of HAVcr-1 codes for a hepatitis A virus cellular receptor. J. Virol. 1998; 72:6621-8.

Ichimura T, Bonventre JV, Bailly V, et al. Kidney injury molecule-1 (KIM-1), a putative epithelial cell adhesion molecule containing a novel immunoglobulin domain, is up-regulated in renal cells after injury. J. Biol. Chem.1998; 273:4135-42.

Meyers JH, Sabatos CA, Chakravarti S, et al. The TIM family regulates autoimmune and allergic diseases. Trends Mol. Med. 2005; 11:362-9.

Meyers JH, Chakravarti S, Schlesinger D, et al. TIM-4 is the ligand for TIM-1, and the TIM-1-TIM4 interaction regulates T cell proliferation. Nat. Immunol. 2005; 6:455-64.

#### Images



Western blot analysis of TIM-1 in human uterus tissue lysate with TIM-1 antibody at (A) 1 and (B) 2  $\mu$ g/mL.

Immunohistochemistry of TIM-1 in human uterus tissue with TIM-1 antibody at 10  $\mu g/mL$ .



Immunofluorescence of TIM-1 in Human Uterus cells with TIM-1 antibody at 20  $\mu\text{g/mL}.$ 

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.