

# ORAI3 Antibody

Catalog # ASC10529

## Product Information

---

<b>Application</b>	WB, ICC, E
<b>Primary Accession</b>	<a href="#">Q9BRQ5</a>
<b>Other Accession</b>	<a href="#">Q9BRQ5</a> , <a href="#">74732916</a>
<b>Reactivity</b>	Human, Mouse, Rat
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	IgG
<b>Calculated MW</b>	31499
<b>Concentration (mg/ml)</b>	1 mg/mL
<b>Conjugate</b>	Unconjugated
<b>Application Notes</b>	ORAI3 antibody can be used for detection of ORAI3 by Western blot at 1 - 4 $\mu$ g/mL. Antibody can also be used for immunocytochemistry starting at 10 $\mu$ g/mL.

## Additional Information

---

<b>Gene ID</b>	93129
<b>Other Names</b>	ORAI3 Antibody: TMEM142C, TMEM142C, Protein orai-3, Transmembrane protein 142C, ORAI calcium release-activated calcium modulator 3
<b>Target/Specificity</b>	ORAI3; This antibody is predicted to have no cross-reactivity to ORAI1 or ORAI2.
<b>Reconstitution &amp; Storage</b>	ORAI3 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.
<b>Precautions</b>	ORAI3 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	ORAI3
<b>Synonyms</b>	TMEM142C
<b>Function</b>	Pore-forming subunit of two major inward rectifying Ca(2+) channels at the plasma membrane: Ca(2+) release-activated Ca(2+) (CRAC) channels and arachidonate-regulated Ca(2+)-selective (ARC) channels (PubMed: <a href="#">16807233</a> , PubMed: <a href="#">17442569</a> , PubMed: <a href="#">19182790</a> , PubMed: <a href="#">19622606</a> , PubMed: <a href="#">19706554</a> , PubMed: <a href="#">20354224</a> , PubMed: <a href="#">32415068</a> ). Assembles with ORAI1 and ORAI2 to form hexameric CRAC channels that mediate Ca(2+)

influx upon depletion of endoplasmic reticulum Ca(2+) store and channel activation by Ca(2+) sensor STIM1, a process known as store-operated Ca(2+) entry (SOCE). Various pore subunit combinations may account for distinct CRAC channel spatiotemporal and cell-type specific dynamics. ORAI1 mainly contributes to the generation of Ca(2+) plateaus involved in sustained Ca(2+) entry and is dispensable for cytosolic Ca(2+) oscillations, whereas ORAI2 and ORAI3 generate oscillatory patterns. CRAC channels assemble in Ca(2+) signaling microdomains where Ca(2+) influx is coupled to calmodulin and calcineurin signaling and activation of NFAT transcription factors recruited to ORAI1 via AKAP5. CRAC channels are the main pathway for Ca(2+) influx in T cells and promote the immune response to pathogens by activating NFAT-dependent cytokine and chemokine transcription (PubMed:[16807233](#), PubMed:[17442569](#), PubMed:[19182790](#), PubMed:[19706554](#), PubMed:[20354224](#), PubMed:[32415068](#)). Assembles with ORAI1 to form channels that mediate store-independent Ca(2+) influx in response to inflammatory metabolites arachidonate or its derivative leukotriene C4, termed ARC and LRC channels respectively (PubMed:[19622606](#), PubMed:[32415068](#)).

#### Cellular Location

Cell membrane; Multi-pass membrane protein. Note=Colocalizes with STIM1 upon store depletion.

#### Tissue Location

Expressed in both naive and effector T helper cells with higher levels in effector cells.

## Background

---

ORAI3 Antibody: Antigen stimulation of immune cells triggers Ca<sup>++</sup> entry through Ca<sup>++</sup> release-activated Ca<sup>++</sup> (CRAC) channels. ORAI3 is one of two mammalian homologs to ORAI1, a recently identified four-transmembrane spanning protein that is an essential component of CRAC. All three homologs have been shown to function as Ca<sup>++</sup> plasma membrane channels gated through interactions with STIM1, the store-activated endoplasmic reticulum Ca<sup>++</sup> sensor. However, ORAI3 channels failed to produce detectable Ca<sup>++</sup> selective currents in cells co-transfected with ORAI3 and STIM1, indicating that ORAI3 channels undergo a lesser degree of depotentiation than ORAI1 or ORAI2. Na<sup>+</sup> currents through ORAI1, 2 and 3 channels were equally inhibited by extracellular Ca<sup>++</sup>, indicating that each have similar affinities for Ca<sup>++</sup> within the selectivity filter.

## References



---

- Lewis RS. Calcium signaling mechanisms in T lymphocytes. *Annu. Rev. Immunol.* 2001; 19:497-521.
- Feske S, Gwack Y, Prakriya M, et al. A mutation in Orai1 causes immune deficiency by abrogating CRAC channel function. *Nature* 2006; 441:179-85.
- Soboloff J, Spassova MA, Dziadek MA, et al. Calcium signals mediated by STIM and Orai proteins - a new paradigm in inter-organelle communication. *Biochim. Biophys. Acta.* 2006; 1763:1161-8.
- Mercer JC, DeHaven WI, Smyth JT, et al. Large store-operated calcium selective currents due to co-expression of Orai1 or Orai2 with the intracellular calcium sensor, Stim1. *J. Biol. Chem.* 2006; 281:24979-90.

## Images

---

Western blot analysis of ORAI3 in A20 cell lysate with ORAI3 antibody at (A) 1, (B) 2 and (C) 4 µg/mL.



Immunocytochemistry of ORAI3 in A20 cells with ORAI3 antibody at 10  $\mu\text{g/mL}$ .

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.