

# TEM2 Antibody

Catalog # ASC10597

## Product Information

---

<b>Application</b>	WB, IF, E, IHC-P
<b>Primary Accession</b>	<a href="#">Q96D21</a>
<b>Other Accession</b>	<a href="#">Q96D21</a> , <a href="#">21362868</a>
<b>Reactivity</b>	Human, Mouse, Rat
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	IgG
<b>Calculated MW</b>	30366
<b>Concentration (mg/ml)</b>	1 mg/mL
<b>Conjugate</b>	Unconjugated
<b>Application Notes</b>	TEM2 antibody can be used for detection of TEM2 by Western blot at 1 - 2 $\mu$ g/mL. Antibody can also be used for immunohistochemistry starting at 2.5 $\mu$ g/mL. For immunofluorescence start at 20 $\mu$ g/mL.

## Additional Information

---

<b>Gene ID</b>	23551
<b>Other Names</b>	GTP-binding protein Rhes, Ras homolog enriched in striatum, Tumor endothelial marker 2, RASD2, TEM2
<b>Target/Specificity</b>	RASD2;
<b>Reconstitution &amp; Storage</b>	TEM2 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.
<b>Precautions</b>	TEM2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	RASD2
<b>Synonyms</b>	TEM2
<b>Function</b>	GTPase signaling protein that binds to and hydrolyzes GTP. Regulates signaling pathways involving G-proteins-coupled receptor and heterotrimeric proteins such as GNB1, GNB2 and GNB3. May be involved in selected striatal competencies, mainly locomotor activity and motor coordination.
<b>Cellular Location</b>	Cell membrane; Lipid-anchor


## Background

TEM2 Antibody: Rhes, also known as tumor endothelial marker 2 (TEM2), is a small GTP-binding protein that is predominantly expressed in the striatal region of the brain. This protein belongs to the RASD subfamily of Ras-related GTP-binding protein superfamily and is thought to play a role in the normal development and function of the brain as mice lacking this gene showed increased anxiety levels and motor coordination deficits. Rhes was identified as TEM2 through analysis of genes whose expression was upregulated in tumor endothelium. Tumor endothelial markers are significantly up-regulated during angiogenesis and neoangiogenesis that are crucial for the growth of solid tumors. TEMs localized on the cell surface and conserved across species are of particular interest for future development of anti-angiogenic therapies.


## References

Vargiu P, Morte B, Manzano J, et al. Thyroid hormone regulation of rhes, a novel RAS homolog gene expressed in the striatum. *Brain Res. Mol. Brain Res.* 2001; 94:1-8.  
Spano D, Branchi I, Rosica A, et al. Rhes is involved in striatal function. *Mol. Cell. Biol.* 2004; 24:5788-96.  
Carson-Walter EB, Watkins DN, Nanda A, et al. Cell surface tumor endothelial markers are conserved in mice and humans. *Cancer Res.* 2001; 61:6649-55.  
Yamamoto Y, Irie K, Nanda A, et al. Direct binding of the human homologue of the Drosophila disc large tumor suppressor gene to seven-pass transmembrane proteins, tumor endothelial marker 5 (TEM5), and a novel TEM5-like protein. *Oncogene* 2004; 23:3889-97.

## Images




Western blot analysis of TEM2 in human colon tissue lysate with TEM2 antibody at (A) 1 and (B) 2 µg/mL.



Immunohistochemistry of TEM2 in human colon tissue with TEM2 antibody at 2.5 µg/mL.

Immunofluorescence of TEM2 in Human Colon cells with TEM2 antibody at 20 µg/mL.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.