

Syntaphilin Antibody

Catalog # ASC10698

Product Information

Application	WB, IF, E, IHC-P
Primary Accession	O15079
Other Accession	O15079 , 21362912
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	53537
Concentration (mg/ml)	1 mg/mL
Conjugate	Unconjugated
Application Notes	Syntaphilin antibody can be used for detection of Syntaphilin by Western blot at 2 - 4 μ g/mL. Despite its predicted molecular weight, Syntaphilin usually migrates at higher molecular weight in SDS-PAGE. Antibody can also be used for immunohistochemistry starting at 5 μ g/mL. For immunofluorescence start at 20 μ g/mL.

Additional Information

Gene ID	9751
Other Names	Syntaphilin, SNPH, KIAA0374
Target/Specificity	SNPH;
Reconstitution & Storage	Syntaphilin antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.
Precautions	Syntaphilin Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	SNPH (HGNC:15931)
Synonyms	KIAA0374
Function	Inhibits SNARE complex formation by absorbing free STX1A.
Cellular Location	Membrane; Single-pass membrane protein. Synapse, synaptosome
Tissue Location	Brain specific. Found in synapses.


Background

Syntaphilin Antibody: Syntaphilin was initially identified in a yeast two-hybrid screen with the carboxy terminal region of Syntaxin-1 as bait. Syntaxin-1 is a key component of the synaptic vesicle docking machinery that forms the SNARE complex with synaptobrevin and SNAP-25. Syntaphilin competes with SNAP-25 for binding to syntaxin-1 and inhibits the formation of the SNARE complex, thereby potentially regulating synaptic vesicle exocytosis. Syntaphilin also binds dynamin-1 and inhibits dynamin-dependent endocytosis. Mice lacking syntaphilin show an increased level of mitochondrial motility and a reduced density of axonal mitochondria. This correlates with an enhanced short-term facilitation and significant impairments in motor ability, suggesting syntaphilin plays a major role in presynaptic function. Multiple isoforms are known to exist.


References

- Lao G, Scheuss V, Gerwin CM, et al. Syntaphilin: a syntaxin-1 clamp that controls SNARE assembly. *Neuron* 2000; 25:191-201.
- Sorensen JB. SNARE complexes prepare for membrane fusion. *Trends Neurosci.* 2005; 28:453-5.
- Das S, Gerwin C, and Sheng ZH. Syntaphilin binds to dynamin-1 and inhibits dynamin-dependent endocytosis. *J. Biol. Chem.* 2003; 278:41221-6.
- Kang J-S, Tian J-H, Pan P-Y, et al. Docking of axonal mitochondria by syntaphilin controls their mobility and affects short-term facilitation. *Cell* 2008; 132:137-148.

Images




Western blot analysis of Syntaphilin in human brain tissue lysate with Syntaphilin antibody at (A) 2 and (B) 4 $\mu\text{g/mL}$.



Immunohistochemistry of Syntaphilin in human brain with Syntaphilin antibody at 5 $\mu\text{g/mL}$.

Immunofluorescence of Syntaphilin in Human Brain cells with Syntaphilin antibody at 20 $\mu\text{g/mL}$.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.