

gp120 Antibody

Catalog # ASC10709

Product Information

Application	WB, E
Primary Accession	Q6BBS3
Other Accession	AAT67551 , 49617711
Reactivity	Virus
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Concentration (mg/ml)	1 mg/mL
Conjugate	Unconjugated
Application Notes	Gp120 antibody can be used for detection of gp120 by Western blot at 0.5 - 1 μ g/mL.

Additional Information

Gene ID	155971
Target/Specificity	env;
Reconstitution & Storage	gp120 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.
Precautions	gp120 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Background

gp120 Antibody: Human immunodeficiency virus type 1 (HIV-1) entry into target cells is directed by the envelope (Env) glycoproteins, which are present on the surface of HIV-1 virion or infected cells in the form of trimers consisting of gp120/gp41 complexes. The surface subunit, gp120, initiates the entry process by interacting sequentially with the CD4 receptor and a co-receptor CCR5 or CXCR4, thereby inducing a conformational change that allows the transmembrane (TM) gp41 subunit to mediate fusion between viral and target cell membranes. Cleavage of Env into its gp120 and gp41 components is necessary for activation of its fusogenic activity.


References

Pinter A. Roles of HIV-1 Env variable regions in viral neutralization and vaccine development. Curr. HIV

Res.2007; 5:542-53.

Alkhatib G and Berger EA. HIV coreceptors: from discovery and designation to new paradigms and promise.
Eur. J. Med. Res.2007; 12:375-84.

Images



Western blot analysis of 1 μ g of gp120 with gp120 antibody at (A) 0.5 and (B) 1 μ g/mL.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.