

BMP15 Antibody

Catalog # ASC10739

Product Information

Application	E, IHC-P
Primary Accession	O95972
Other Accession	NP_005439 , 257743454
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	45055
Concentration (mg/ml)	1 mg/mL
Conjugate	Unconjugated
Application Notes	BMP15 antibody can be used for detection of BMP15 by Western blot at 1 - 2 μ g/mL.

Additional Information

Gene ID	9210
Other Names	Bone morphogenetic protein 15, BMP-15, Growth/differentiation factor 9B, GDF-9B, BMP15, GDF9B
Target/Specificity	BMP15;
Reconstitution & Storage	BMP15 antibody can be stored at 4°C for three months and -20°C, stable for up to one year.
Precautions	BMP15 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	BMP15
Synonyms	GDF9B
Function	May be involved in follicular development. Oocyte-specific growth/differentiation factor that stimulates folliculogenesis and granulosa cell (GC) growth.
Cellular Location	Secreted.


Background

BMP15 Antibody: BMP15, or GDF9B is a member of the bone morphogenetic protein family which is part of the transforming growth factor-beta superfamily that are involved in embryonic development and adult tissue homeostasis. BMP15 is expressed exclusively in the oocyte. It is an oocyte-specific growth/differentiation factor that may be involved in oocyte maturation and follicular development. Defects in BMP15 are the cause of ovarian dysgenesis type 2. BMP15 could be used as an oogenesis marker to track human amniotic fluid stem cells differentiation into the oocyte-like cells.

References

Massague J. The transforming growth factor-beta family. *Ann. Rev. Cell. Biol.* 1990; 6:597-641
Aaltonen J, Laitinen MP, Vuojolainen K, et al. Human growth differentiation factor 9 (GDF-9) and its novel homolog GDF-9B are expressed in oocytes during early folliculogenesis. *J. Clin. Endocrinol. Metab.* 1999; 84:2744-50.
Laitinen M, Vuojolainen K, Jaatinen R, et al. A novel growth differentiation factor-9 (GDF-9) related factor is coexpressed with GDF-9 in mouse oocytes during folliculogenesis. *Mech. Dev.* 1998; 78:135-40.
Galloway SM, Grogan SM, Wilson T et al. Bmp15 mutations and ovarian function. *Mol. Cell Endocrinol.* 2002; 191:15-8

Images



Immunohistochemistry of BMP15 in human prostate tissue with BMP15 antibody at 10 µg/mL

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.