

IL-34 Antibody

Catalog # ASC10756

Product Information

Application	WB, E
Primary Accession	Q6ZMJ4
Other Accession	ACD13474 , 187610438
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	27482
Concentration (mg/ml)	1 mg/mL
Conjugate	Unconjugated
Application Notes	IL-34 antibody can be used for the detection of IL-34 by Western blot at 0.25 µg/mL. Despite its predicted molecular weight, IL-34 will often run at higher molecular weight in SDS-PAGE.

Additional Information

Gene ID	146433
Other Names	Interleukin-34, IL-34, IL34, C16orf77
Target/Specificity	IL34;
Reconstitution & Storage	IL-34 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.
Precautions	IL-34 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	IL34
Synonyms	C16orf77
Function	Cytokine that promotes the proliferation, survival and differentiation of monocytes and macrophages. Promotes the release of pro-inflammatory chemokines, and thereby plays an important role in innate immunity and in inflammatory processes. Plays an important role in the regulation of osteoclast proliferation and differentiation, and in the regulation of bone resorption. Signaling via CSF1R and its downstream effectors stimulates phosphorylation of MAPK1/ERK2 AND MAPK3/ERK1.

Cellular Location	Secreted.
Tissue Location	Detected in the sinusoidal epithelium in the red pulp of spleen (at protein level). Predominantly expressed in spleen Also detected in a range of other tissues including heart, brain, lung, liver, kidney, thymus, testis, ovary, small intestine, prostate and colon.


Background

IL-34 Antibody: Interleukin-34 (IL-34) was identified through a functional screening of a library of secreted proteins after transfection of an embryonic kidney cell line with recombinant cDNAs. Flow cytometry and monocyte viability assays demonstrated increased survival and/or proliferation of monocytes induced by IL-34 and specific binding of IL-34 to CD14⁺ monocytes. IL-34 is a second ligand to the colony stimulating factor (CSF)-1 receptor in addition to CSF1. Like CSF1, IL-34 stimulates phosphorylation of ERK1 and ERK2 in human monocytes and induces differentiation of bone marrow cells into macrophages.

References

Lin H, Lee E, Hestir K, et al. Discovery of a cytokine and its receptor by functional screening of the extracellular proteome. *Science* 2008; 320:807-11.

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.