

KLRA2 Antibody

Catalog # ASC10848

Product Information

Application	WB, IF, E, IHC-P
Primary Accession	Q60660
Other Accession	Q60660 , 2497636
Reactivity	Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	33607
Concentration (mg/ml)	1 mg/mL
Conjugate	Unconjugated
Application Notes	KLRA2 antibody can be used for detection of KLRA2 by Western blot at 0.5 - 1 μ g/mL. Antibody can also be used for immunohistochemistry starting at 2.5 μ g/mL. For immunofluorescence start at 20 μ g/mL.

Additional Information

Other Names	Killer cell lectin-like receptor 2, Lymphocyte antigen 49b, Ly-49b, T-cell surface glycoprotein Ly-49B, Klra2, Ly49-b, Ly49b
Target/Specificity	Klra2;
Reconstitution & Storage	KLRA2 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.
Precautions	KLRA2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	Klra2
Synonyms	Ly49-b, Ly49b
Function	Receptor on natural killer (NK) cells for class I MHC.
Cellular Location	Membrane; Single-pass type II membrane protein.

Background

KLRA2 Antibody: KLRA2 (also known as Ly49B) is a member of the LY49 family of receptors in Natural Killer (NK) cells that bind to major histocompatibility complex (MHC) class 1. These proteins are classified as either activating or inhibitory receptors based on whether they possess an immunoreceptor tyrosine-based inhibitory motif (ITIM) in their cytoplasmic region (for inhibitory receptors), or an immunoreceptor tyrosine-based activation motif (ITAM) that transmits activating signals resulting in phosphorylation of several substrates. KLRA2 is thought to be an inhibitory receptor and associates with the protein tyrosine phosphatases SHP1, SHP2, and SHIP in transfected cells. This antibody is predicted to not cross-react with other LY49 receptors.

References


Gosselin P, Lusignan Y, Brennan J, et al. The NK2.1 receptor is encoded by Ly-49C and its expression is regulated by MHC class I alleles. *Int. Immunol.* 1997; 533-40.

Smith HR, Karlhofer FM, and Yokoyama WM. Ly-49 multigene family expressed by IL-2-activated NK cells. *J. Immunol.* 1994; 153:1068-79.


Brennan J, Mager D, Jefferies W, et al. Expression of different members of the Ly-49 gene family defines distinct natural killer cell subsets and cell adhesion properties. *J. Exp. Med.* 1994; 180:2287-95.

Saleh A, Davies GE, Pascal V, et al. Identification of probabilistic transcriptional switches in the Ly49 gene cluster: a eukaryotic mechanism for selective gene activation. *Immunity* 2004; 21:55-66.

Images



Western blot analysis of KLRA2 in mouse spleen tissue lysate with KLRA2 antibody at (A) 0.5 and (B) 1 µg/mL.



Immunohistochemistry of KLRA2 in mouse spleen tissue with KLRA2 antibody at 2.5 µg/mL.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.