

# FNIP2 Antibody

Catalog # ASC10854

## Product Information

---

<b>Application</b>	WB, E, IHC-P
<b>Primary Accession</b>	<a href="#">Q9P278</a>
<b>Other Accession</b>	<a href="#">NP_065891</a> , <a href="#">154689769</a>
<b>Reactivity</b>	Human, Mouse, Rat
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	IgG
<b>Calculated MW</b>	122115
<b>Concentration (mg/ml)</b>	1 mg/mL
<b>Conjugate</b>	Unconjugated
<b>Application Notes</b>	FNIP2 antibody can be used for detection of FNIP2 by Western blot at 1 - 2 µg/mL. Antibody can also be used for immunohistochemistry starting at 5 µg/mL.

## Additional Information

---

<b>Gene ID</b>	57600
<b>Other Names</b>	Folliculin-interacting protein 2, FNIP1-like protein, O6-methylguanine-induced apoptosis 1 protein, FNIP2, FNIP1, KIAA1450, MAPO1
<b>Target/Specificity</b>	FNIP2; Multiple isoforms of FNIP2 are known to exist. This antibody is predicted to not cross-react with FNIP1.
<b>Reconstitution &amp; Storage</b>	FNIP2 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.
<b>Precautions</b>	FNIP2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	FNIP2 {ECO:0000303   PubMed:18663353, ECO:0000312   HGNC:HGNC:29280}
<b>Function</b>	Binding partner of the GTPase-activating protein FLCN: involved in the cellular response to amino acid availability by regulating the non-canonical mTORC1 signaling cascade controlling the MiT/TFE factors TFE3 (PubMed: <a href="#">18663353</a> , PubMed: <a href="#">31672913</a> , PubMed: <a href="#">36103527</a> ). Required to promote FLCN recruitment to lysosomes and interaction with Rag GTPases, leading to activation of the non- canonical mTORC1 signaling (By similarity). In low-amino acid conditions, component of the lysosomal folliculin complex (LFC) on the membrane of lysosomes, which inhibits the GTPase-activating

activity of FLCN, thereby inactivating mTORC1 and promoting nuclear translocation of TFEB and TFE3 (PubMed:[31672913](#), PubMed:[36103527](#)). Upon amino acid restimulation, disassembly of the LFC complex liberates the GTPase- activating activity of FLCN, leading to activation of mTORC1 and subsequent inactivation of TFEB and TFE3 (PubMed:[31672913](#)). Together with FLCN, regulates autophagy: following phosphorylation by ULK1, interacts with GABARAP and promotes autophagy (PubMed:[25126726](#)). In addition to its role in mTORC1 signaling, also acts as a co-chaperone of HSP90AA1/Hsp90: inhibits the ATPase activity of HSP90AA1/Hsp90, leading to activate both kinase and non-kinase client proteins of HSP90AA1/Hsp90 (PubMed:[18403135](#)). Acts as a scaffold to load client protein FLCN onto HSP90AA1/Hsp90 (PubMed:[18403135](#)). Competes with the activating co-chaperone AHSA1 for binding to HSP90AA1, thereby providing a reciprocal regulatory mechanism for chaperoning of client proteins (PubMed:[18403135](#)). May play a role in the signal transduction pathway of apoptosis induced by O6-methylguanine-mispaired lesions (By similarity).

#### Cellular Location

Lysosome membrane. Cytoplasm. Note=Colocalizes with FLCN in the cytoplasm.

#### Tissue Location

Widely expressed with highest levels in muscle, nasal mucosa, salivary gland, uvula, fat, liver, heart, placenta and pancreas (PubMed:[18403135](#), PubMed:[18663353](#), PubMed:[27353360](#)) Moderately expressed in the lung, small intestine, kidney and brain Lower levels detected in renal cell carcinoma than in normal kidney tissue (PubMed:[18403135](#)). Higher levels detected in oncocytoma tumors than in normal kidney. Higher levels detected in renal cell carcinoma tumors than in normal kidney tissue (PubMed:[27353360](#))

## Background

---

FNIP2 Antibody: FNIP2 is the second protein found to interact with folliculin, the product of the Birt-Hogg-Dube (BHD) gene. Folliculin is thought to act as a tumor suppressor as mutations or loss of heterozygosity in this gene are associated with BHD syndrome-related renal tumors. Folliculin and FNIP1, a protein that shares 49% identity to FNIP2, bind to AMPK, an important energy sensor in cells that negatively regulates the mammalian target of rapamycin (mTOR), a protein that is thought to be the master switch for cell growth and proliferation. FNIP1 and FNIP2 are able to form homo- and heteromeric multimers, suggesting these proteins may have a functional relationship.

## References

---

Hasumi H, Baba M, Hong S-B, et al. Identification and characterization of a novel folliculin-interacting protein FNIP2. *Gene* 2008; 415:60-7.

Takagi Y, Kobayashi T, Shiono M, et al. Interaction of folliculin (Birt-Hogg-Dube gene product) with novel Fnip1-like (FnipL/Fnip2) protein. *Oncogene* 2008; 27:5339-47.



Vocke CD, Yang Y, Pavlovich CP, et al. High frequency of somatic frameshift BHD mutations in Birt-Hogg-Dube-associated renal tumors. *J. Natl. Cancer Inst.* 2005; 97:931-5.

Baba M, Hong SB, Sharma M, et al. Folliculin encoded by the BHD gene interacts with a binding protein, FNIP1, and AMPK, and is involved in AMPK and mTOR signaling. *Proc. Natl. Acad. Sci. USA* 2006; 103:15552-7.

## Images

---

Western blot analysis of FNIP2 in rat skeletal muscle lysate with FNIP2 antibody at (A) 1 and (B) 2 µg/mL.



Immunohistochemistry of FNIP2 in mouse skeletal muscle tissue with FNIP2 antibody at 5  $\mu$ g/mL.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.