

ZBED3 Antibody

Catalog # ASC10885

Product Information

Application	WB, E
Primary Accession	Q96IU2
Other Accession	NP_115743 , 14150185
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	25132
Concentration (mg/ml)	1 mg/mL
Conjugate	Unconjugated
Application Notes	ZBED3 antibody can be used for detection of ZBED3 by Western blot at 1 - 2 μ g/mL. Despite its predicted molecular weight, ZBED3 often migrates at a higher position in SDS-PAGE.

Additional Information

Gene ID	84327
Other Names	Zinc finger BED domain-containing protein 3, Axin-interacting protein, ZBED3
Target/Specificity	ZBED3; This antibody is predicted to have no cross-reactivity to ZBED1 or ZBED2.
Reconstitution & Storage	ZBED3 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.
Precautions	ZBED3 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	ZBED3
Function	Acts as a positive regulator in the activation of the canonical Wnt/beta-catenin signaling pathway by stabilizing cytoplasmic beta-catenin (By similarity). Involved in transcription activation of Wnt target gene expression (By similarity). Plays a role in symmetric division of blastomeres in the early stages of embryogenesis via regulation of mitotic spindle central positioning and organization of the F-actin filament network (By similarity).
Cellular Location	Cytoplasm {ECO:0000250 UniProtKB:Q9D0L1}. Membrane {ECO:0000250 UniProtKB:Q9D0L1}. Secreted

Tissue Location

Secreted in blood plasma, and expressed in skeletal muscle and adipose tissue (at protein level)


Background

ZBED3 Antibody: Axin is a key modulator of the Wnt/beta-catenin pathway and acts as a scaffold protein in phosphorylating and degrading cytoplasmic beta-catenin. ZBED3, a member of the Zinc finger BED-type domain containing superfamily, has been recently identified as an Axin-interacting protein and has been suggested to modulate the Wnt/beta-catenin signaling pathway. RNA interference (RNAi) experiments targeting ZBED3 RNA attenuated Wnt-induced beta-catenin accumulation, lymphoid enhancer binding factor-1-dependent luciferase reporter activity, and Wnt gene expression.

References

Chen T, Li M, Ding Y, et al. Identification of Zinc-finger BED domain containing 3 (Zbed3) as a novel axin-interacting protein that activates Wnt/b-catenin signaling. J. Biol. Chem.2009; 284:6683-9.

Images



Western blot analysis of ZBED3 in A20 cell lysate with ZBED3 antibody at 1 µg/mL.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.