

Fibulin 3 Antibody

Catalog # ASC10920

Product Information

Application WB, IF, ICC, E
Primary Accession Q12805

Other Accession NP_001034437, 9665262
Reactivity Human, Mouse, Rat

Host Rabbit
Clonality Polyclonal
Isotype IgG
Calculated MW 54641
Concentration (mg/ml) 1 mg/mL
Conjugate Unconjugated

Application Notes Fibulin 3 antibody can be used for detection of Fibulin 3 by Western blot at

0.5 - 1 [g/mL. Antibody can also be used for immunocytochemistry starting at

20 g/mL. For immunofluorescence start at 20 g/mL.

Additional Information

Gene ID 2202

Other Names EGF-containing fibulin-like extracellular matrix protein 1, Extracellular protein

S1-5, Fibrillin-like protein, Fibulin-3, FIBL-3, EFEMP1, FBLN3, FBNL

Target/Specificity EFEMP1; At least three isoforms of Fibulin 3 are known to exist. This antibody

is predicted to not cross-react with other Fibulin proteins.

Reconstitution & Storage Fibulin 3 antibody can be stored at 4°C for three months and -20°C, stable for

up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high

temperatures.

Precautions Fibulin 3 Antibody is for research use only and not for use in diagnostic or

therapeutic procedures.

Protein Information

Name EFEMP1

Synonyms FBLN3, FBNL

Function Binds EGFR, the EGF receptor, inducing EGFR autophosphorylation and the

activation of downstream signaling pathways. May play a role in cell adhesion

and migration. May function as a negative regulator of chondrocyte

differentiation. In the olfactory epithelium, it may regulate glial cell migration,

differentiation and the ability of glial cells to support neuronal neurite

outgrowth.

Cellular Location Secreted, extracellular space, extracellular matrix. Note=Localizes to the

lamina propria underneath the olfactory epithelium

{ECO:0000250 | UniProtKB:O35568}

Tissue Location In the eye, associated with photoreceptor outer and inner segment regions,

the nerve fiber layer, outer nuclear layer and inner and outer plexiform layers

of the retina

Background

Fibulin 3 Antibody: Fibulin 3, also known as epidermal growth factor (EGF)-containing fibulin-like extracellular matrix protein 1 (EFEMP1), is a member of the fibulin family of extracellular glycoproteins, a group of proteins that are widely distributed and frequently associated with vascular and elastic tissues. The fibulin proteins typically contain a tandem array of EGF-like repeats and a fibulin-type COOH-terminal module. Aberrant accumulation of Fibulin 3 in the endoplasmic reticulum of retinal pigment epithelial cells has been shown to be associated with inherited forms of macular degeneration, but the loss of Fibulin 3 expression does not lead to macular degeneration but rather the appearance of hernias due to a reduction of elastic fibers of fascial connective tissue. Recent experiments have shown that expression of Fibulin 3 promotes tumor growth and may thus be a therapeutic target.


References

Kobayashi N, Kostka G, Garbe JH, et al. A comparative analysis of the fibulin protein family. Biochemical characterization, binding interactions, and tissue localization. J. Biol. Chem. 2007; 282:11805-16. Marmorstein LY, Munier FL, Arsenijevic Y, et al. Aberrant accumulation of EFEMP1 underlies drusen formation in Malattia Leventinese and age-related macular degeneration. Proc. Natl. Acad. Sci. USA 2002; 99:13067-72.

McLaughlin PJ, Bakall B, Choi J, et al. Lack of fibulin-3 causes early aging and herniation, but not macular degeneration in mice. Hum. Mol. Genet. 2007; 16:3059-70.

Seeliger H, Camaj P, Ischenko I, et al. EFEMP1 expression promotes in vivo tumor growth in human pancreatic adenocarcinoma. Mol. Cancer Res. 2009; 7:189-98.


Images



Western blot analysis of Fibulin 3 in HeLa cell lysate with Fibulin 3 antibody at (A) 0.5 and (B) 1 μ g/mL.



Immunocytochemistry of Fibulin 3 in HeLa cells with Fibulin 3 antibody at 20 µg/mL.



Immunofluorescence of Fibulin 3 in HeLa cells with Fibulin 3 antibody at 20 $\mu g/mL$.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.