

PLEKHM2 Antibody

Catalog # ASC11016

Product Information

Application	WB, IF, E, IHC-P
Primary Accession	Q8IWE5
Other Accession	CAH72018 , 55666400
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	112780
Concentration (mg/ml)	1 mg/mL
Conjugate	Unconjugated
Application Notes	PLEKHM2 antibody can be used for detection of PLEKHM2 by Western blot at 0.5 µg/mL. Antibody can also be used for immunohistochemistry starting at 5 µg/mL. For immunofluorescence start at 20 µg/mL.

Additional Information

Gene ID	23207
Other Names	Pleckstrin homology domain-containing family M member 2, PH domain-containing family M member 2, Salmonella-induced filaments A and kinesin-interacting protein, SifA and kinesin-interacting protein, PLEKHM2, KIAA0842, SKIP
Target/Specificity	PLEKHM2;
Reconstitution & Storage	PLEKHM2 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.
Precautions	PLEKHM2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	PLEKHM2 (HGNC:29131)
Function	Plays a role in lysosomes movement and localization at the cell periphery acting as an effector of ARL8B. Required for ARL8B to exert its effects on lysosome location, recruits kinesin-1 to lysosomes and hence direct their movement toward microtubule plus ends. Binding to ARL8B provides a link from lysosomal membranes to plus-end-directed motility (PubMed: 22172677 , PubMed: 24088571 , PubMed: 25898167 , PubMed: 28325809). Critical factor involved in NK cell-mediated cytotoxicity. Drives the polarization of cytolytic

granules and microtubule-organizing centers (MTOCs) toward the immune synapse between effector NK lymphocytes and target cells (PubMed:[24088571](#)). Required for maintenance of the Golgi apparatus organization (PubMed:[22172677](#)). May play a role in membrane tubulation (PubMed:[15905402](#)).

Cellular Location

Cytoplasm. Lysosome membrane; Peripheral membrane protein; Cytoplasmic side


Background

PLEKHM2 Antibody: PLEKHM2, also known as SKIP, is a member of the M family of Pleckstrin homology domain-containing proteins. While little is known of PLEKHM2, a recent study examining differential gene expression in human hematopoietic stem cells has shown it to be specifically expressed in stem cells, suggesting that PLEKHM2 may play a role in erythroid commitment and development. Other studies have shown that PLEKHM2 is required for interaction with the Salmonella virulence factor SifA for Salmonella pathogenesis, suggesting that PLEKHM2 has cellular roles other than in the developing embryo.


References

Liu X-L, Yuan J-Y, Zhang J-W, et al. Differential gene expression in human hematopoietic stem cells specified toward erythroid, megakaryotic, and granulocytic lineage. *J. Leuk. Biol.* 2007; 82:986-1002.
Diacovich L, Dumont A, Lafitte D, et al. Interaction between the SifA virulence factor and its host target SKIP is essential for Salmonella pathogenesis. *J. Biol. Chem.* 2009; 284:33151-60.

Images




Western blot analysis of PLEKHM2 in rat brain tissue lysate with PLEKHM2 antibody at 0.5 µg/mL in (A) the absence and (B) the presence of blocking peptide.



Immunohistochemistry of PLEKHM2 in human liver tissue with FAM59A antibody at 5 µg/mL.

Immunofluorescence of PLEKHM2 in Human Brain cells with PLEKHM2 antibody at 20 µg/mL.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.