

SIRT4 Antibody

Catalog # ASC11137

Product Information

Application	WB, E
Primary Accession	Q9Y6E7
Other Accession	EAW98182 , 119618588
Reactivity	Human, Mouse, Rat
Host	Chicken
Clonality	Polyclonal
Isotype	IgY
Calculated MW	35188
Concentration (mg/ml)	1 mg/mL
Conjugate	Unconjugated
Application Notes	SIRT4 antibody can be used for detection of SIRT4 by Western blot at 1 - 2 μ g/mL.

Additional Information

Gene ID	23409
Other Names	NAD-dependent protein deacetylase sirtuin-4 {ECO:0000255 HAMAP-Rule:MF_03161}, 3.5.1.- {ECO:0000255 HAMAP-Rule:MF_03161}, NAD-dependent ADP-ribosyltransferase sirtuin-4 {ECO:0000255 HAMAP-Rule:MF_03161}, 2.4.2.- {ECO:0000255 HAMAP-Rule:MF_03161}, Regulatory protein SIR2 homolog 4 {ECO:0000255 HAMAP-Rule:MF_03161}, SIR2-like protein 4 {ECO:0000255 HAMAP-Rule:MF_03161}, SIRT4 {ECO:0000255 HAMAP-Rule:MF_03161}, SIR2L4
Target/Specificity	SIRT4;
Reconstitution & Storage	SIRT4 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.
Precautions	SIRT4 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	SIRT4 {ECO:0000255 HAMAP-Rule:MF_03161, ECO:0000312 HGNC:HGNC:14932}
Function	Acts as a NAD-dependent protein lipoamidase, biotinylase, deacetylase and ADP-ribosyl transferase (PubMed: 16959573 , PubMed: 17715127 , PubMed: 24052263 , PubMed: 25525879). Catalyzes more efficiently removal of

lipoyl- and biotinyl- than acetyl-lysine modifications (PubMed:[24052263](#), PubMed:[25525879](#)). Inhibits the pyruvate dehydrogenase complex (PDH) activity via the enzymatic hydrolysis of the lipoamide cofactor from the E2 component, DLAT, in a phosphorylation-independent manner (PubMed:[25525879](#)). Catalyzes the transfer of ADP-ribosyl groups onto target proteins, including mitochondrial GLUD1, inhibiting GLUD1 enzyme activity (PubMed:[16959573](#), PubMed:[17715127](#)). Acts as a negative regulator of mitochondrial glutamine metabolism by mediating mono ADP-ribosylation of GLUD1: expressed in response to DNA damage and negatively regulates anaplerosis by inhibiting GLUD1, leading to block metabolism of glutamine into tricarboxylic acid cycle and promoting cell cycle arrest (PubMed:[16959573](#), PubMed:[17715127](#)). In response to mTORC1 signal, SIRT4 expression is repressed, promoting anaplerosis and cell proliferation (PubMed:[23663782](#)). Acts as a tumor suppressor (PubMed:[23562301](#), PubMed:[23663782](#)). Also acts as a NAD-dependent protein deacetylase: mediates deacetylation of 'Lys-471' of MLYCD, inhibiting its activity, thereby acting as a regulator of lipid homeostasis (By similarity). Does not seem to deacetylate PC (PubMed:[23438705](#)). Controls fatty acid oxidation by inhibiting PPARA transcriptional activation (PubMed:[24043310](#)). Impairs SIRT1-PPARA interaction probably through the regulation of NAD(+) levels (PubMed:[24043310](#)). Down-regulates insulin secretion (PubMed:[17715127](#)).

Cellular Location

Mitochondrion matrix {ECO:0000255 | HAMAP- Rule:MF_03161, ECO:0000269 | PubMed:16079181, ECO:0000269 | PubMed:16959573, ECO:0000269 | PubMed:17715127}

Tissue Location

Detected in vascular smooth muscle and striated muscle. Detected in insulin-producing beta-cells in pancreas islets of Langerhans (at protein level). Widely expressed. Weakly expressed in leukocytes and fetal thymus.

Background


SIRT4 Antibody: The Silent Information Regulator (SIR2) family of genes are highly conserved from prokaryotes to eukaryotes and have important functions in the regulation of metabolism, growth and differentiation, inflammation, cellular survival, as well as in senescence and lifespan extension. Sirtuins, including SIRT1-7, are human homologs of yeast Sir2p. Sirtuins are NAD⁺-dependent histone/protein deacetylases (HDAC) which regulate cellular metabolism, e.g. energy metabolism, and thereby are associated with aging and several age-related diseases. SIRT4 localizes to mitochondria, inhibits glutamate dehydrogenase, and is thought to be involved in the regulation of insulin secretion.

References

Salminen A. SIRT1: regulation of longevity via autophagy. *Cell Signal*2009; 21:1356-60.
 Afshar G and Murnane JP. Characterization of a human gene with sequence homology to *Saccharomyces cerevisiae* Sir 2. *Gene*1999; 234:161-8.
 Guarente L. Sirtuins as potential targets for metabolic syndrome. *Nature*2006; 444:868-74.
 Vaziri H, Dessain SK, Ng Eaton E, et al. hSIR2 (SIRT1) functions as an NAD-dependent p53 deacetylase. *Cell*2001; 107:149-59.

Images

Western blot analysis of SIRT4 in human liver tissue lysate with SIRT4 antibody at 1 µg/mL.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.