

TMEM192 Antibody

Catalog # ASC11296

Product Information

Application	WB, IF, E, IHC-P
Primary Accession	Q8IY95
Other Accession	NP_001093859 , 154240704
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	30922
Concentration (mg/ml)	1 mg/mL
Conjugate	Unconjugated
Application Notes	TMEM192 antibody can be used for detection of TMEM192 by Western blot at 0.5 µg/mL. Antibody can also be used for immunohistochemistry starting at 5 µg/mL. For immunofluorescence start at 20 µg/mL.

Additional Information

Gene ID	201931
Other Names	Transmembrane protein 192, TMEM192
Target/Specificity	TMEM192; At least two isoforms of TMEM192 are known to exist; this antibody will recognize both. TMEM192 antibody is predicted to not cross-react with other TMEM proteins.
Reconstitution & Storage	TMEM192 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.
Precautions	TMEM192 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	TMEM192
Cellular Location	Lysosome membrane; Multi-pass membrane protein. Late endosome
Tissue Location	Strongly expressed in kidney, liver, lung and pancreas.

Background


TMEM192 Antibody: TMEM192 is a widely expressed, four transmembrane protein that localizes to late endosomal/lysosomal membranes. This localization is mediated by two dileucine motifs present at the amino-terminus of TMEM192. A cysteine residue near the carboxy residue is responsible for disulfide bond formation in TMEM192 homodimers. Little is known about the function of this protein.

References


Schroder B, Wrocklage C, Hasilik A, et al. Molecular characterisation of 'transmembrane protein 192'(TMEM192, a novel protein of the lysosomal membrane. *Biol. Chem.* 2010; 391:695-704.

Zhu H, Liang Z, and Li G. Rabex-5 is a Rab22 effector and mediates a Rab22-Rab5 signaling cascade in endocytosis. *Mol. Biol. Cell* 2009; 20:4720-9.


Images



Western blot analysis of TMEM192 in SK-N-SH cell lysate with TMEM192 antibody at 0.5 µg/mL.



Immunohistochemistry of TMEM192 in mouse brain tissue with TMEM192 antibody at 5 µg/mL.



Immunofluorescence of TMEM192 in mouse brain tissue with TMEM192 antibody at 20 µg/mL.