

HDGFRP2 Antibody

Catalog # ASC11351

Product Information

Application	WB, IF, E, IHC-P
Primary Accession	Q7Z4V5
Other Accession	EAW69215 , 48255931
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	74317
Concentration (mg/ml)	1 mg/mL
Conjugate	Unconjugated
Application Notes	HDGFRP2 antibody can be used for detection of HDGFRP2 by Western blot at 0.5 - 1 μ g/mL. Antibody can also be used for immunohistochemistry starting at 5 μ g/mL. For immunofluorescence start at 20 μ g/mL.

Additional Information

Gene ID	84717
Other Names	Hepatoma-derived growth factor-related protein 2, HRP-2, Hepatoma-derived growth factor 2, HDGF-2, HDGFRP2, HDGF2
Target/Specificity	HDGFRP2; At least two isoforms of HDGFRP2 are known to exist; this antibody will recognize both isoforms. This antibody is predicted to not cross-react with other HDGFRP family members
Reconstitution & Storage	HDGFRP2 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.
Precautions	HDGFRP2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	HDGFL2 (HGNC:14680)
Function	Acts as an epigenetic regulator of myogenesis in cooperation with DPF3a (isoform 2 of DPF3/BAF45C) (PubMed: 32459350). Associates with the BAF complex via its interaction with DPF3a and HDGFL2-DPF3a activate myogenic genes by increasing chromatin accessibility through recruitment of SMARCA4/BRG1/BAF190A (ATPase subunit of the BAF complex) to myogenic gene promoters (PubMed: 32459350). Promotes the repair of DNA double-strand breaks (DSBs) through the homologous recombination

pathway by facilitating the recruitment of the DNA endonuclease RBBP8 to the DSBs (PubMed:[26721387](#)). Preferentially binds to chromatin regions marked by H3K9me3, H3K27me3 and H3K36me2 (PubMed:[26721387](#), PubMed:[32459350](#)). Involved in cellular growth control, through the regulation of cyclin D1 expression (PubMed:[25689719](#)).

Cellular Location

Nucleus. Cytoplasm {ECO:0000250|UniProtKB:Q925G1}

Tissue Location

Widely expressed. High expression is found in heart, skeletal muscle, ovary and testis. Overexpression is frequently observed in hepatocellular carcinoma samples


Background

HDGFRP2 Antibody: Hepatoma-derived growth factor-related protein 2 (HDGFRP2) is a widely expressed member of protein family that contains the HDGF related proteins 1-4 and lens epithelial derived growth factor (LEDGF). These family members show extensive homology at the amino terminus (termed the hath region). Little is known of the role of HDGFRP2, but like the related LEDGF, HDGFRP2 can bind HIV-1 integrase and stimulate HIV-1 integrase activity in vitro.

References



El-Tahir HM, Dietz F, Dringen R, et al. Expression of Hepatoma-derived growth factor family members in the adult central nervous system. *BMC Neurosci.* 2006; 7:6.
Izumoto Y, Kuroda T, Harada H, et al. Hepatoma-derived growth factor belongs to a gene family in mice showing significant homology in the amino terminus. *Biochem. Biophys. Res. Commun.* 1997; 238:26-32
Cherepanov P, Devroe E, Silver PA, et al. Identification of an evolutionarily conserved domain in human lens epithelium-derived growth factor/transcriptional co-activator p75 (LEDGF/p75) that binds HIV-1 integrase. *J. Biol. Chem.* 2004; 279:48883-92.

Images



Western blot analysis of HDGFRP2 in rat lung tissue lysate with HDGFRP2 antibody at 2 µg/mL .

Immunohistochemistry of HDGFRP2 in rat liver tissue with HDGFRP2 antibody at 5 µg/mL.



Immunofluorescence of HDGFRP2 in rat liver tissue with HDGFRP2 antibody at 20 $\mu\text{g/mL}$.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.