

# **UHRF1BP1L** Antibody

Catalog # ASC11368

### **Product Information**

**Application** WB, IF, E, IHC-P

Primary Accession <u>A0JNW5</u>

Other Accession NP\_055869, 55749644
Reactivity Human, Mouse

Host Rabbit
Clonality Polyclonal
Isotype IgG
Calculated MW 164199
Concentration (mg/ml) 1 mg/mL
Conjugate Unconjugated

**Application Notes**UHRF1BP1L antibody can be used for detection of UHRF1BP1L by Western

blot at 1 [g/mL. Antibody can also be used for immunohistochemistry starting

at 2.5 g/mL. For immunofluorescence start at 20 g/mL.

#### **Additional Information**

**Gene ID** 23074

Other Names UHRF1-binding protein 1-like, UHRF1BP1L, KIAA0701

**Target/Specificity** UHRF1BP1L; At least two isoforms of UHRF1BP1L are known to exist; this

antibody will recognize both isoforms. UHRF1BP1L antibody is predicted to

not cross-react with UHRF1BP1

**Reconstitution & Storage** UHRF1BP1L antibody can be stored at 4°C for three months and -20°C, stable

for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged

high temperatures.

**Precautions**UHRF1BP1L Antibody is for research use only and not for use in diagnostic or

therapeutic procedures.

#### **Protein Information**

Name BLTP3B ( HGNC:29102)

**Function** Tube-forming lipid transport protein which mediates the transfer of lipids

between membranes at organelle contact sites (PubMed:<u>35499567</u>). Required for retrograde traffic of vesicle clusters in the early endocytic pathway to the

Golgi complex (PubMed:20163565, PubMed:35499567).

**Cellular Location** Cytoplasm, cytosol. Early endosome. Note=Localizes on a subpopulation of

vesicle clusters in the early endocytic pathway

## **Background**

UHRF1BP1L Antibody: The Ubiquitin-like containing PHD and RING finger domains 1-binding protein 1-like (UHRF1BP1L) is closely related to UHRF1BP1, also known as ICBP90, a transcription and cell cycle regulator that specifically binds to the histone H3 N-terminal tail. While little is known of UHRF1BP1L, UHRF1BP1 is required for proper heterochromatin formation in mammalian cells. Furthermore, UHRF1BP1 is thought to be a pivotal target for the ERK1/2 signaling pathway to control the proliferation of Jurkat T cells, suggesting that UHRF1BP1L may also be involved in chromatin regulation and cell proliferation.

#### References


Hopfner R, Mousli M, Jeltsch JM, et al. ICBP90, a novel human CCAAT binding protein, involved in the regulation of Topoisomerase IIα expression. Cancer Res. 2000; 60:121-8

Unoki M, Nishidate T and Nakamura Y. ICBP90, an E2F-1 target, recruits HDAC1 and binds to methyl-CpG through its SRA domain. Oncogene 2004; 23:7601-10


Karagianni P, Amazit L, Qin J, et al. ICBP90, a novel methyl K9 H3 binding protein linking protein ubiquitination with heterochromatin formation. Mol. Cell Biol. 2008; 28:705-17

Fang Z, Xing F, Bronner C, et al. ICBP90 mediates the ERK1/2 signaling to regulate the proliferation of Jurkat T cells. Cell Immunol. 2009; 257:80-7

## **Images**




Western blot analysis of UHRF1BP1L in mouse brain tissue lysate with UHRF1BP1L antibody at 1 µg/mL



Immunohistochemistry of UHRF1BP1L in human kidney tissue with UHRF1BP1L antibody at 2.5  $\mu$ g/mL.

Immunofluorescence of UHRF1BP1L in human kidney tissue with UHRF1BP1L antibody at 20 µg/mL.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.