

Periphilin Antibody

Catalog # ASC11436

Product Information

Application	WB, IF, E, IHC-P
Primary Accession	Q8NEY8
Other Accession	NP_958848 , 48255929
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	52737
Concentration (mg/ml)	1 mg/mL
Conjugate	Unconjugated
Application Notes	Periphilin antibody can be used for detection of Periphilin by Western blot at 1 µg/mL. Antibody can also be used for immunohistochemistry starting at 2.5 µg/mL. For immunofluorescence start at 5 µg/mL.

Additional Information

Gene ID	51535
Other Names	Periphilin-1, Gastric cancer antigen Ga50, PPHLN1
Target/Specificity	PPHLN1;
Reconstitution & Storage	Periphilin antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.
Precautions	Periphilin Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	PPHLN1 (HGNC:19369)
Function	Component of the HUSH complex, a multiprotein complex that mediates epigenetic repression. The HUSH complex is recruited to genomic loci rich in H3K9me3 and is probably required to maintain transcriptional silencing by promoting recruitment of SETDB1, a histone methyltransferase that mediates further deposition of H3K9me3. In the HUSH complex, contributes to the maintenance of the complex at chromatin (PubMed: 26022416). Acts as a transcriptional corepressor and regulates the cell cycle, probably via the HUSH complex (PubMed: 15474462 , PubMed: 17963697). The HUSH complex is also involved in the silencing of unintegrated retroviral DNA: some part of the retroviral DNA formed immediately after infection remains unintegrated in

the host genome and is transcriptionally repressed (PubMed:[30487602](#)). May be involved in epithelial differentiation by contributing to epidermal integrity and barrier formation (PubMed:[12853457](#)).

Cellular Location

Nucleus. Cytoplasm. Chromosome. Note=In undifferentiated keratinocytes expressed in speckle-type nuclear granules and at the nuclear membrane, but in the differentiated keratinocytes colocalized with periplakin at the cell periphery and at cell-cell junctions (PubMed:12853457) Localizes to chromatin (PubMed:26022416).

Tissue Location

Ubiquitous..


Background

Periphilin Antibody:Periphilin, known as PPHLN1 or gastric cancer antigen Ga50, plays a role in epithelial differentiation and contributes to epidermal integrity and barrier formation. It interacts with periplakin, a known precursor of the cornified cell envelope.Periphilin 1 is ubiquitously expressed and localizes to nucleus and cytoplasm. Existing as eight alternatively spliced isoforms,Periphilin is highly insoluble and contains a putative nuclear localization signal (NLS) within its N-terminal half, a prerequisite for the formation of insoluble complexes and a possible caspase recognition sequence and a potential nuclear export signal.Periphilin may play an important role in the nervous system.

References



Kazerounian S and Aho S. Characterization of periphilin, a widespread, highly insoluble nuclear protein and potential constituent of the keratinocyte cornified envelope. *J. Biol. Chem.* 2003; 278:36707-17.
Kurita M, Suzuki H, Masai H, et al. Overexpression of CR/periphilin downregulates Cdc7 expression and induces S-phase arrest. *Biochem. Biophys. Res. Commun.* 2004; 324:554-61
Soehn AS, Pham TT, Schaeferhoff K, et al.Periphilin is strongly expressed in the murine nervous system and is indispensable for murine development. *Genesis* 2009; 47:697-707
Soehn AS, Franck T, Biskup S, et al.Periphilin is a novel interactor of synphilin-1, a protein implicated in Parkinson's disease. *Neurogenetics* 2010; 11:203-15

Images



Western blot analysis ofPeriphilin in mouse colon tissue lysate withPeriphilin antibody at 1 µg/mL in (A) the absence and (B) the presence of blocking peptide.

Immunohistochemistry ofPeriphilin in human colon tissue withPeriphilin antibody at 2.5 µg/mL.



Immunofluorescence of Periphilin in human colon tissue with Periphilin antibody at 5 $\mu\text{g/mL}$.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.