10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

# SELK Antibody

Catalog # ASC11524

## Product Information

| | |
|---|---|
| **Application** | WB, IF, E |
| **Primary Accession** | Q9Y6D0 |
| **Other Accession** | NP_067060, 25014099 |
| **Reactivity** | Human, Mouse, Rat |
| **Host** | Rabbit |
| **Clonality** | Polyclonal |
| **Isotype** | IgG |
| **Calculated MW** | 10645 |
| **Concentration (mg/ml)** | 1 mg/mL |
| **Conjugate** | Unconjugated |
| **Application Notes** | SELK antibody can be used for detection of SELK by Western blot at 1 - 2 g/mL. For immunofluorescence start at 20 g/mL. |

## Additional Information

| | |
|---|---|
| **Gene ID** | 58515 |
| **Other Names** | Selenoprotein K, SelK, SELK |
| **Target/Specificity** | SELK; |
| **Reconstitution & Storage** | SELK antibody can be stored at 4℃ for three months and -20℃, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures. |
| **Precautions** | SELK Antibody is for research use only and not for use in diagnostic or therapeutic procedures. |

## Protein Information

| | |
|---|---|
| **Name** | SELENOK {ECO:0000303\|PubMed:27645994, ECO:0000312\|HGNC:HGNC:30394} |
| **Function** | Required for Ca(2+) flux in immune cells and plays a role in T-cell proliferation and in T-cell and neutrophil migration (By similarity). Involved in endoplasmic reticulum-associated degradation (ERAD) of soluble glycosylated proteins (PubMed:22016385). Required for palmitoylation and cell surface expression of CD36 and involved in macrophage uptake of low-density lipoprotein and in foam cell formation (By similarity). Together with ZDHHC6, required for palmitoylation of ITPR1 in immune cells, leading to regulate ITPR1 stability and function (PubMed:25368151). Plays a role in protection of cells from ER stress- induced apoptosis (PubMed:20692228). Protects cells from oxidative stress when overexpressed in cardiomyocytes |

(PubMed:16962588).

| | |
|---|---|
| **Cellular Location** | Endoplasmic reticulum membrane; Single-pass membrane protein. Cell membrane; Single-pass membrane protein. Note=Probably mainly localized in the ER |
| **Tissue Location** | Highly expressed in heart. |


## Background

SELK Antibody: SELK is a selenoprotein, containing a selenocysteine residue at its active site. SELK is localized to the endoplasmic reticulum and is highly expressed in the heart, where it may function as an antioxidant. SELK can bind to proteins such as the ER-associated degradation (ERAD) components p97 ATPase, Derlin-1 and -2, and SelS as well as proteins in the oligosaccharyltransferase complex, suggesting that SELK may be involved in the Derlin-dependent ERAD of glycosylated misfolded proteins. SELK is also thought to be important in Ca2+ flux in immune cells, and is cleaved by m-calpain in resting macrophages. When these macrophages are activated through different Toll-like receptors, this cleavage is inhibited


## References

Lu C, Qiu F, Zhou H, et al. Identification and characterization of selenoprotein K: An antioxidant in cardiomyocytes. FEBS Lett. 2006; 580:5189-97
Shchedrina VA, Everly RA, Zhang Y, et al. Selenoprotein K binds multiprotein complexes and is involved in the regulation of endoplasmic homeostasis. J. Biol. Chem. 2011; 286:42937-48.
Huang Z, Hoffmann FW, Norton RL, et al. Selenoprotein K is a novel target of m-calpain, and cleavage is regulated by Toll-like receptor-induced calpastatin in macrophages. J. Biol. Chem. 2011; 286:34830-8.

## Images




Western blot analysis of SELK in A20 cell lysate with SELK antibody at (A) 1 and (B) 2 µg/mL.



Immunofluorescence of SELK in human heart tissue with SELK antibody at 20 µg/mL.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.