

TSPY1S Antibody

Catalog # ASC11583

Product Information

Application	WB, E
Primary Accession	Q01534
Other Accession	NP_001184171 , 308522758
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	35012
Concentration (mg/ml)	1 mg/mL
Conjugate	Unconjugated
Application Notes	TSPY1S antibody can be used for detection of TSPY1S by Western blot at 1 - 2 μ g/mL.

Additional Information

Gene ID	100289087;7258;728137
Other Names	Testis-specific Y-encoded protein 1, Cancer/testis antigen 78, CT78, TSPY1, TSPY
Target/Specificity	TSPY1; TSPY1S antibody is human and mouse reactive. At least three isoforms of TSPY1 are known to exist; this antibody will detect only TSPY1S
Reconstitution & Storage	TSPY1S antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.
Precautions	TSPY1S Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	TSPY1
Synonyms	TSPY
Function	May be involved in sperm differentiation and proliferation.
Cellular Location	Cytoplasm. Nucleus. Note=Predominantly cytoplasmic. Also found in nucleus
Tissue Location	Specifically expressed in testicular tissues. Isoform 1 and isoform 2 are expressed in spermatogonia and spermatocytes. Found in early testicular carcinoma in situ, spermatogonial cells in testicular tissues of 46,X,Y female

and in prostate cancer cell lines.

Background

TSPY1S Antibody: Testis-specific protein on Y chromosome (TSPY1) is an ampliconic gene on the Y chromosome that has been associated with gonadoblastoma. Recent experiments have shown that in androgen-dependent testicular germ-cell tumors, TSPY1 can repress the androgen-bound androgen receptor (AR), a member of the nuclear steroid hormone receptor family that acts as a ligand-inducible transcription factor, suggesting that TSPY1 is a repressor of cell proliferation in germ-cell tumors and potentially in normal gonadal cells during early development. Two distinct isoforms of TSPY1, TSPY1L and TSPY1S, are known to exist.

References


Arnemann J, Jakubiczka S, Thuring S, et al. Cloning and sequence analysis of a human Y-chromosome-derived, testicular cDNA, TSPY. *Genomics* 1991; 11:108-114

Lau YF. Gonadoblastoma, testicular and prostate cancers, and the TSPY gene. *Am. J. Hum. Genet.* 1999; 64:921-7.

Akimoto C, Ueda T, Inoue K, et al. Testis-specific protein on Y chromosome (TSPY) represses the activity of the androgen receptor in androgen-dependent testicular germ-cell tumors. *Proc. Natl. Acad. Sci. USA* 2010; 107:19891-6.

Krick R, Jakubiczka S, and Arnemann J. Expression, alternative splicing and haplotype analysis of transcribed testis specific protein (TSPY) genes. *Gene* 2003; 302:11-9.

Images



Western blot analysis of TSPY1S in A20 cell lysate with TSPY1S antibody at 1 µg/mL in (A) the absence and (B) the presence of blocking peptide.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.