

# IL-15 Antibody

Catalog # ASC11615

### **Product Information**

**Application** WB, E, IHC-P **Primary Accession** P40933

Other Accession NP\_000576, 10835153
Reactivity Human, Mouse, Rat

Host Rabbit
Clonality Polyclonal
Isotype IgG
Calculated MW 18086
Concentration (mg/ml) 1 mg/mL
Conjugate Unconjugated

**Application Notes** IL-15 antibody can be used for detection of IL-15 by Western blot at 1 - 2

□g/mL.

## **Additional Information**

Gene ID 3600

Other Names Interleukin-15, IL-15, IL15

**Target/Specificity** IL15; At least two isoforms of IL-15 are known to exist.

**Reconstitution & Storage** IL-15 antibody can be stored at 4°C for three months and -20°C, stable for up

to one year.

**Precautions** IL-15 Antibody is for research use only and not for use in diagnostic or

therapeutic procedures.

## **Protein Information**

Name IL15

**Function** Cytokine that plays a major role in the development of inflammatory and

protective immune responses to microbial invaders and parasites by modulating immune cells of both the innate and adaptive immune systems (PubMed:15123770). Stimulates the proliferation of natural killer cells, T-cells

and B-cells and promotes the secretion of several cytokines

(PubMed:<u>8178155</u>, PubMed:<u>9326248</u>). In monocytes, induces the production of IL8 and monocyte chemotactic protein 1/CCL2, two chemokines that attract

neutrophils and monocytes respectively to sites of infection

(PubMed:<u>9326248</u>). Unlike most cytokines, which are secreted in soluble form, IL15 is expressed in association with its high affinity IL15RA on the surface of IL15-producing cells and delivers signals to target cells that express IL2RB and IL2RG receptor subunits (PubMed:<u>10233906</u>, PubMed:<u>23104097</u>, PubMed:<u>8026467</u>). Binding to its receptor triggers the phosphorylation of

JAK1 and JAK3 and the recruitment and subsequent phosphorylation of signal transducer and activator of transcription-3/STAT3 and STAT5 (PubMed:7568001). In mast cells, induces the rapid tyrosine phosphorylation of STAT6 and thereby controls mast cell survival and release of cytokines such as IL4 (By similarity).

**Cellular Location** [Isoform IL15-S48AA]: Secreted.

**Tissue Location**Most abundant in placenta and skeletal muscle. It is also detected in the

heart, lung, liver and kidney. IL15-S21AA is preferentially expressed in tissues

such as testis and thymus

# **Background**

IL-15 Antibody: Interleukin 15 (IL-15) is a cytokine that regulates T and natural killer cell activation and proliferation. This cytokine and IL-2 share many biological activities as both have been found to bind common hematopoietin receptor subunits, and may compete for the same receptor, and thus negatively regulate each other's activity. The number of CD8+ memory cells is shown to be controlled by a balance between IL-15 and IL-2. This cytokine induces the activation of JAK kinases, as well as the phosphorylation and activation of transcription activators STAT3, STAT5, and STAT6. In mouse, studies suggest that IL-15 may increase the expression of apoptosis inhibitor Bcl-xL, possibly through the transcription activation activity of STAT6, and thus prevent apoptosis.

#### References


Bodnar A, Nizsaloczki E, Mocsar G, et al. A biophysical approach to IL-2 and IL-15 receptor function: localization, conformation and interactions. Immunol. Lett. 2008; 116:117-25.

Ku CC, Murakami M, Sakamoto A, et al. Control of homeostasis of CD8+ memory T cells by opposing cytokines. Science 2000; 288:675-8.

Waldmann T, Tagaya Y, and Bamford R. Interleukin-2, interleukin-15, and their receptors. Int. Rev. Immunol. 1998; 16:205-26.


Masuda A, Matsuguchi T, Yamaki K, et al. Interleukin-15 prevents mouse mast cell apoptosis through STAT6-mediate Bcl-xL expression. J. Biol. Chem. 2001; 276:26107-13.

# **Images**



Western blot analysis of IL-15 in rat spleen tissue lysate with IL-15 antibody at (A) 1 and (B) 2 µg/mL

Immunohistochemistry of IL-15 in spleen tissue with IL-15 antibody at 2.5  $\mu$ g/ml.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.