

TLX3 Antibody

Catalog # ASC11663

Product Information

Application	WB, IF, E, IHC-P
Primary Accession	O43711
Other Accession	NP_066305 , 50355978
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	31867
Concentration (mg/ml)	1 mg/mL
Conjugate	Unconjugated
Application Notes	TLX3 antibody can be used for detection of TLX3 by Western blot at 0.5 - 1 μ g/mL.

Additional Information

Gene ID	30012
Other Names	T-cell leukemia homeobox protein 3, Homeobox protein Hox-11L2, TLX3, HOX11L2
Target/Specificity	TLX3; TLX3 antibody is predicted to not cross-react with other TLX proteins.
Reconstitution & Storage	TLX3 antibody can be stored at 4°C for three months and -20°C, stable for up to one year.
Precautions	TLX3 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	TLX3
Synonyms	HOX11L2
Cellular Location	Nucleus.

Background

TLX3 Antibody: The Hox proteins play a role in patterns of embryonic development and cellular differentiation by regulating downstream target genes. TLX3, also known as homeobox11-like1 (HOX11L2), is a DNA-binding nuclear transcription factor that is expressed in a subset of the primary sensory nervous system. TLX3, along with the related TLX1, is a selector transcription factor that promotes an excitatory

glutamatergic neuronal phenotype over an inhibitory GABAergic phenotype opposing LBX1 signals during dorsal spinal cord development. Chromosomal translocations of the TLX3 gene have been shown to result in some forms of T-cell acute lymphoblastic leukemia (T-ALL).

References


Dear TN, Sanchez-Garcia I, and Rabbitts TH, et al. The Hox11 gene encodes a DNA-binding nuclear transcription factor belonging to a distinct family of homeobox genes. *Proc. Natl. Acad. Sci. USA* 1993; 90:4431-5.

Patterson KD and Kried PA. Hox11-family genes XHox11 and XHox11L2 in xenopus: XHoxL2 expression is restricted to a subset of the primary sensory neurons. *Dev. Dyn.* 1999; 214:34-43.


Cheng L, Arata A, Mizuguchi R, et al. Tlx3 and Tlx1 are post-mitotic selector genes determining the glutamatergic over GABAergic cell fates. *Nat. Neurosci.* 2004; 7:510-7.

Cheng L, Samad OA, Xu Y, et al. Lbx1 and Tlx3 are opposing switches in determining GABAergic versus glutamatergic transmitter phenotypes. *Nat. Neurosci.* 2005; 8:1510-5.


Images



Western blot analysis of TLX3 in human spleen tissue lysate with TLX3 antibody at (A) 0.5 and (B) 1 µg/mL.



Immunohistochemistry of TLX3 in human spleen tissue with TLX3 antibody at 2.5 µg/ml.



Immunofluorescence of TLX3 in human spleen tissue with TLX3 antibody at 20 µg/ml.