

CD160 Antibody

Catalog # ASC11698

Product Information

Application	WB, IF, E
Primary Accession	O95971
Other Accession	NP_008984 , 5901910
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	19810
Concentration (mg/ml)	1 mg/mL
Conjugate	Unconjugated
Application Notes	CD160 antibody can be used for detection of CD160 by Western blot at 1 - 2 μ g/ml.

Additional Information

Gene ID	11126
Other Names	CD160 antigen, Natural killer cell receptor BY55, CD160, CD160, BY55
Target/Specificity	CD160; CD160 antibody is human and mouse reactive. Multiple isoforms of CD160 are known to exist.
Reconstitution & Storage	CD160 antibody can be stored at 4°C for three months and -20°C, stable for up to one year.
Precautions	CD160 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	CD160 {ECO:0000303 PubMed:16809620, ECO:0000312 HGNC:HGNC:17013}
Function	[CD160 antigen]: Receptor on immune cells capable to deliver stimulatory or inhibitory signals that regulate cell activation and differentiation. Exists as a GPI-anchored and as a transmembrane form, each likely initiating distinct signaling pathways via phosphoinositol 3-kinase in activated NK cells and via LCK and CD247/CD3 zeta chain in activated T cells (PubMed: 11978774 , PubMed: 17307798 , PubMed: 19109136). Receptor for both classical and non-classical MHC class I molecules (PubMed: 12486241 , PubMed: 9973372). In the context of acute viral infection, recognizes HLA-C and triggers NK cell cytotoxic activity, likely playing a role in anti-viral innate immune response (PubMed: 12486241). On CD8+ T cells, binds HLA-A2-B2M in complex with a viral peptide and provides a costimulatory signal to activated/memory T cells

(PubMed:[9973372](#)). Upon persistent antigen stimulation, such as occurs during chronic viral infection, may progressively inhibit TCR signaling in memory CD8+ T cells, contributing to T cell exhaustion (PubMed:[25255144](#)). On endothelial cells, recognizes HLA-G and controls angiogenesis in immune privileged sites (PubMed:[16809620](#)). Receptor or ligand for TNF superfamily member TNFRSF14, participating in bidirectional cell-cell contact signaling between antigen presenting cells and lymphocytes. Upon ligation of TNFRSF14, provides stimulatory signal to NK cells enhancing IFNG production and anti-tumor immune response (By similarity). On activated CD4+ T cells, interacts with TNFRSF14 and down-regulates CD28 costimulatory signaling, restricting memory and alloantigen-specific immune response (PubMed:[18193050](#)). In the context of bacterial infection, acts as a ligand for TNFRSF14 on epithelial cells, triggering the production of antimicrobial proteins and pro-inflammatory cytokines (By similarity).

Cellular Location

[CD160 antigen]: Cell membrane; Lipid-anchor, GPI-anchor

Tissue Location

Expression is restricted to functional NK and cytotoxic T lymphocytes. Expressed in viral-specific effector memory and terminally differentiated effector memory CD8+ T cells. Expressed in memory and activated CD4+ T cell subsets (at protein level) (PubMed:11978774, PubMed:18193050, PubMed:25255144, PubMed:9743336) Expressed at high levels in intraepithelial lymphocytes (at protein level) (PubMed:9743336). Expressed in both alpha-beta and gamma-delta CD8+ T cell subsets (at protein level) (PubMed:11978774, PubMed:18193050, PubMed:9743336). Expressed in umbilical vein endothelial cells (at protein level) (PubMed:16809620). Expressed in monocytes and at lower levels in B cells (PubMed:23761635). Isoform 3: Expressed exclusively in activated NK cells (at protein level) (PubMed:19109136).

Background



CD160, also known as BY55, is a lipid-anchored cell membrane glycoprotein that contains one immunoglobulin-like domain (1). It is expressed in small intestine, spleen and functional NK and T cytotoxic lymphocytes (1,2). CD160 exists as a disulfide-linked homomultimer that functions as a receptor for MHC (major histocompatibility complex) molecules and is thought to regulate the function of NK cells (2,3). Additionally, CD160 interacts with TNFRSF14 and, via this interaction, is able to negatively regulate CD4+ T cell activation, indicating a role in immune system regulation (4).

References

- Anumanthan A, Bensussan A, Boumsell L, et al. Cloning of BY55, a novel Ig superfamily member expressed on NK cells, CTL, and intestinal intraepithelial lymphocytes. *J. Immunol.* 1998; 161:2780-90.
- Agrawal S, Marquet J, Freeman GJ, et al. Cutting edge: MHC class I triggering by a novel cell surface ligand costimulates proliferation of activated human T cells. *J. Immunol.* 1999; 162:1223-6.
- Le Bouteiller P, Tabiasco J, Polgar B, et al. CD160: a unique activating NK cell receptor. *Immunol. Lett.* 2011;138:93-6
- Chabot S, Jabrane-Ferrat N, Bigot K, et al. A novel antiangiogenic and vascular normalization therapy targeted against human CD160 receptor. *J. Exp. Med.* 2011; 208:973-86.

Images

Western blot analysis of CD160 in K562 cell lysate with CD160 antibody at 1 µg/ml in (A) the absence and (B) the presence of blocking peptide.



Immunofluorescence of CD160 in Jurkat cells tissue with CD160 antibody at 5 $\mu\text{g/mL}$.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.