

# GRK6 Antibody

Catalog # ASC11746

## Product Information

---

<b>Application</b>	WB, IF, E, IHC-P
<b>Primary Accession</b>	<a href="#">P43250</a>
<b>Other Accession</b>	<a href="#">NP_001004106</a> , <a href="#">51896039</a>
<b>Reactivity</b>	Human, Mouse, Rat
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	IgG
<b>Calculated MW</b>	65991
<b>Concentration (mg/ml)</b>	1 mg/mL
<b>Conjugate</b>	Unconjugated
<b>Application Notes</b>	GRK6 antibody can be used for detection of GRK6 by Western blot at 1 - 2 $\mu$ g/ml. Antibody can also be used for Immunohistochemistry starting at 5 $\mu$ g/mL. For immunofluorescence start at 20 $\mu$ g/mL.

## Additional Information

---

<b>Gene ID</b>	2870
<b>Other Names</b>	G protein-coupled receptor kinase 6, 2.7.11.16, G protein-coupled receptor kinase GRK6, GRK6, GPRK6
<b>Target/Specificity</b>	GRK6; GRK6 antibody is human, mouse and rat reactive. Multiple isoforms of GRK6 are known to exist. This antibody is predicted to not cross-react with other members of the GRK protein family.
<b>Reconstitution &amp; Storage</b>	GRK6 antibody can be stored at 4°C for three months and -20°C, stable for up to one year.
<b>Precautions</b>	GRK6 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	GRK6
<b>Synonyms</b>	GPRK6
<b>Function</b>	Specifically phosphorylates the activated forms of G protein- coupled receptors. Such receptor phosphorylation initiates beta- arrestin-mediated receptor desensitization, internalization, and signaling events leading to their desensitization. Seems to be involved in the desensitization of D2-like dopamine receptors in striatum and chemokine receptor CXCR4 which is critical for CXCL12-induced cell chemotaxis (By similarity). Phosphorylates rhodopsin (RHO) (in vitro) and a non G-protein-coupled receptor: LRP6 during

Wnt signaling (in vitro).

**Cellular Location** Membrane; Lipid-anchor.

**Tissue Location** Widely expressed..

## Background

---

The G protein-coupled receptor kinases (GRKs) are a versatile family of kinases that play a critical role in cancer metastasis through their regulation of G-protein coupled receptors (GPCRs) involved in growth factor mediated cell migration (1,2). Phosphorylation of receptors by GRKs appears to be strictly dependent on the receptor being in its agonist-activated state (1). GRK6 is one of 7 members of the GRK serine/threonine kinase subfamily, which has been shown to modulate the Wnt signaling pathway via phosphorylation of LRP6 (3,4), and the insulin-like growth factor signaling pathway (4). GRK6 may also play a role in immune system function (5).

## References

---

Inglese J, Freedman NJ, Koch WJ, et al. Structure and mechanism of the G protein-coupled receptor kinases. *J. Biol. Chem.* 1993; 268:23735-8.


Raghuwanshi SK, Smith N, Rivers EJ, et al. G protein-coupled receptor kinase 6 deficiency promotes angiogenesis, tumor progression, and metastasis. *J. Immunol.* 2013; 190:5329-36.

Benovic JL and Gomez J. Molecular cloning and expression of GRK6. A new member of the G protein-coupled receptor kinase family. *J. Biol. Chem.* 1993; 268:19521-7.


Chen M, Philipp M, Wang J, et al. G Protein-coupled receptor kinases phosphorylate LRP6 in the Wnt pathway. *J. Biol. Chem.* 2009; 284:35040-8.

## Images

---




Western blot analysis of GRK6 in rat small intestine tissue lysate with GRK6 antibody at 1 µg/ml.



Immunohistochemistry of GRK6 in human small intestine tissue with GRK6 antibody at 5 µg/mL.

Immunofluorescence of GRK6 in human small intestine tissue with GRK6 antibody at 20 µg/mL.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.