

# ME1 Antibody

Catalog # ASC11773

## Product Information

---

<b>Application</b>	WB, E, IHC-P
<b>Primary Accession</b>	<a href="#">P48163</a>
<b>Other Accession</b>	<a href="#">NP_002386</a> , <a href="#">4505143</a>
<b>Reactivity</b>	Human, Mouse, Rat
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	IgG
<b>Calculated MW</b>	64150
<b>Concentration (mg/ml)</b>	1 mg/mL
<b>Conjugate</b>	Unconjugated
<b>Application Notes</b>	ME1 antibody can be used for detection of ME1 by Western blot at 1 - 2 $\mu$ g/ml. Antibody can also be used for Immunohistochemistry at 5 $\mu$ g/mL.

## Additional Information

---

<b>Gene ID</b>	4199
<b>Other Names</b>	NADP-dependent malic enzyme, NADP-ME, 1.1.1.40, Malic enzyme 1, ME1
<b>Target/Specificity</b>	ME1; ME1 antibody is human, mouse and rat reactive. ME1 antibody is predicted not to cross-react with ME2.
<b>Reconstitution &amp; Storage</b>	ME1 antibody can be stored at 4°C for three months and -20°C, stable for up to one year.
<b>Precautions</b>	ME1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	ME1 ( <a href="#">HGNC:6983</a> )
<b>Function</b>	Catalyzes the oxidative decarboxylation of (S)-malate in the presence of NADP(+) and divalent metal ions, and decarboxylation of oxaloacetate.
<b>Cellular Location</b>	Cytoplasm.
<b>Tissue Location</b>	Expressed in all tissues tested including liver, placenta and white adipose tissue.

## Background

---

ME1, also known as NADP-ME, MES or HUMNDME, may play an important role as a housekeeping protein within the cell (1). ME1 is a 572 amino acid cytoplasmic protein that belongs to the malic enzyme family (2). It is expressed ubiquitously with highest expression in liver and white adipose tissue. ME1 functions as an NADP-dependent enzyme that generates NADPH for fatty acid biosynthesis (3). The activity of this enzyme, the reversible oxidative decarboxylation of malate, links the glycolytic and citric acid cycles (3,4). ME1 is regulated by both thyroid hormone levels and the amount of carbohydrates in the diet (5).

## References

---

Loeber G, Dworkin MB, Infante A, et al. Characterization of cytosolic malic enzyme in human tumor cells. *FEBS Lett.* 1994; 344:181-6.


Gonzalez-Manchon C, Ferrer M, Ayuso MS, et al. Cloning, sequencing and functional expression of a cDNA encoding a NADP-dependent malic enzyme from human liver. *Gene* 1995;159:255-60.

Gonzalez-Manchon C, Butta N, Ferrer M, et al. Molecular cloning and functional characterization of the human cytosolic malic enzyme promoter: thyroid hormone responsiveness. *DNA Cell Biol.* 1997;16:533-44.


Yang Z, Lanks CW and Tong L. Molecular mechanism for the regulation of human mitochondrial NAD(P)<sup>+</sup>-dependent malic enzyme by ATP and fumarate. *Structure* 2002; 10:951-60.

## Images

---



Western blot analysis of ME1 in 293 cell lysate with ME1 antibody at 1 µg/ml.



Immunohistochemistry of ME1 in mouse liver tissue with ME1 antibody at 5 µg/mL.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.