

WNT5 Antibody

Catalog # ASC11881

Product Information

Application	WB, E, IHC-P
Primary Accession	Q9H1J7
Other Accession	NP_110402 , 17402919
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	40323
Concentration (mg/ml)	1 mg/mL
Conjugate	Unconjugated
Application Notes	WNT5 antibody can be used for detection of WNT5 by Western blot at 1 - 2 μ g/ml. For immunofluorescence start at 20 μ g/mL.

Additional Information

Gene ID	81029
Other Names	Protein Wnt-5b, WNT5B
Target/Specificity	WNT5; WNT5 antibody is human, mouse and rat reactive. WNT5 antibody will detect both WNT5A and WNT5B. At least two isoforms of WNT5B are known to exist; this antibody will detect both isoforms.
Reconstitution & Storage	WNT5 antibody can be stored at 4°C for three months and -20°C, stable for up to one year.
Precautions	WNT5 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	WNT5B
Function	Ligand for members of the frizzled family of seven transmembrane receptors. Probable developmental protein. May be a signaling molecule which affects the development of discrete regions of tissues. Is likely to signal over only few cell diameters (By similarity).
Cellular Location	Secreted, extracellular space, extracellular matrix

Background

WNT5A and WNT5B are near-identical members of the WNT gene family, which consists of structurally related genes which encode secreted signaling proteins that have been implicated in oncogenesis and in several developmental processes, including regulation of cell fate and patterning during embryogenesis (1,2). WNT5A and WNT5B coordinate chondrocyte proliferation and differentiation by differentially regulating cyclin D1 and p130 expression, as well as chondrocyte-specific Col2a1 expression (3). Up-regulation of WNT5A and WNT5B has been observed in several types of human cancers, suggesting that they may be useful as diagnostic markers or potent therapeutic targets (4).

References


Saitoh T and Katoh M. Molecular cloning and characterization of human WNT5B on chromosome 12p13.3 region. *Int. J. Oncol.* 2001; 19:347-51.

Serio RN. Wnt of the two horizons: putting stem cell self-renewal and cell fate determination into context. *Stem Cells Dev.* 2014; epub Apr. 24.


Yang Y, Topol L, Lee H, et al. Wnt5a and Wnt5b exhibit distinct activities in coordinating chondrocyte proliferation and differentiation. *Development* 2003; 130:1003-15.

Saitoh T and Katoh M. Expression and regulation of WNT5A and WNT5B in human cancer: up-regulation of WNT5A by TNFalpha in MKN45 cells and up-regulation of WNT5B by beta-estradiol in MCF-7 cells. *Int. J. Mol. Med.* 2002; 10:345-9.

Images



Western blot analysis of WNT5 in Jurkat cell lysate with WNT5 antibody at 1 µg/ml in (A) the absence and (B) the presence of blocking peptide.



Immunohistochemistry of WNT5 in rat colon tissue with WNT5 antibody at 5 µg/mL.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.