

CX3CR1 Antibody (biotin)

Chemokines & Cytokines, Cancer, Infectious Disease

Catalog # ASC12208

Product Information

Application	E
Primary Accession	P49238
Other Accession	NP_001328
Reactivity	Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Clone Names	CX3CR1
Calculated MW	40396
Concentration (mg/ml)	1 mg/mL
Conjugate	biotin

Additional Information

Gene ID	1524
Alias Symbol	CX3CR1
Other Names	CX3CR1 Antibody: V28, CCRL1, GPR13, CMKDR1, GPRV28, CMKBRL1, CX3C chemokine receptor 1, Beta chemokine receptor-like 1, C-X3-C CKR-1
Reconstitution & Storage	CX3CR1 antibody can be stored at 4 °C for three months and -20 °C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.
Precautions	CX3CR1 Antibody (biotin) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	CX3CR1 {ECO:0000303 PubMed:12551893, ECO:0000312 HGNC:HGNC:2558}
Function	Receptor for the C-X3-C chemokine fractalkine (CX3CL1) present on many early leukocyte cells; CX3CR1-CX3CL1 signaling exerts distinct functions in different tissue compartments, such as immune response, inflammation, cell adhesion and chemotaxis (PubMed: 12055230 , PubMed: 23125415 , PubMed: 9390561 , PubMed: 9782118). CX3CR1-CX3CL1 signaling mediates cell migratory functions (By similarity). Responsible for the recruitment of natural killer (NK) cells to inflamed tissues (By similarity). Acts as a regulator of inflammation process leading to atherogenesis by mediating macrophage and monocyte recruitment to inflamed atherosclerotic plaques, promoting cell

survival (By similarity). Involved in airway inflammation by promoting interleukin 2-producing T helper (Th2) cell survival in inflamed lung (By similarity). Involved in the migration of circulating monocytes to non-inflamed tissues, where they differentiate into macrophages and dendritic cells (By similarity). Acts as a negative regulator of angiogenesis, probably by promoting macrophage chemotaxis (PubMed:[14581400](#), PubMed:[18971423](#)). Plays a key role in brain microglia by regulating inflammatory response in the central nervous system (CNS) and regulating synapse maturation (By similarity). Required to restrain the microglial inflammatory response in the CNS and the resulting parenchymal damage in response to pathological stimuli (By similarity). Involved in brain development by participating in synaptic pruning, a natural process during which brain microglia eliminates extra synapses during postnatal development (By similarity). Synaptic pruning by microglia is required to promote the maturation of circuit connectivity during brain development (By similarity). Acts as an important regulator of the gut microbiota by controlling immunity to intestinal bacteria and fungi (By similarity). Expressed in lamina propria dendritic cells in the small intestine, which form transepithelial dendrites capable of taking up bacteria in order to provide defense against pathogenic bacteria (By similarity). Required to initiate innate and adaptive immune responses against dissemination of commensal fungi (mycobiota) component of the gut: expressed in mononuclear phagocytes (MNP) and acts by promoting induction of antifungal IgG antibodies response to confer protection against disseminated *C.albicans* or *C.auris* infection (PubMed:[29326275](#)). Also acts as a receptor for C-C motif chemokine CCL26, inducing cell chemotaxis (PubMed:[20974991](#)).

Cellular Location

Cell membrane; Multi-pass membrane protein

Tissue Location

Expressed in lymphoid and neural tissues (PubMed:7590284). Expressed in lymphocyte subsets, such as natural killer (NK) cells, gamma-delta T-cells and terminally differentiated CD8(+) T-cells (PubMed:12055230). Expressed in smooth muscle cells in atherosclerotic plaques (PubMed:14581400)

Background


CX3CR1 Antibody: CX3CR1 is one of the chemokine receptors that are required as coreceptors for HIV infection. The genes encoding human, murine, and rat CX3CR1 were cloned and designated V28 and CMKBRL1, CX3CR1, and RBS11, respectively. The encoded seven transmembrane protein was recently identified as the receptor for a novel transmembrane molecule, fractalkine, and renamed CX3CR1. Recently, CX3CR1 was found to serve as a coreceptor for HIV-1 and HIV-2 envelope fusion and virus infection, which can be inhibited by fractokine. CX3CR1 mediates leukocyte migration and adhesion. CX3CR1 is expressed in a variety of human tissues and cell lines.

References

Raport et al. Gene 1995;163:295-9.
Combadiere et al. DNA Cell Biol 1995;14:673-80.
Combadiere et al. Biochem Biophys Res Commun 1998;253:728-32.
Harrison et al. Neurosci Lett 1994;169:85-9.

Images

Figure 1 ELISA Validation



Coating Antigen: immunogen peptide, 2093P, 10 µg/mL, incubate at 4 °C overnight. Detection Antibodies: CX3CR1, ASC12208, dilution: 0.5-1000 ng/mL, incubate at RT for 1 hr. ASC12208 was detected by HRP-conjugated streptavidin at 1:5,000 was detected by anti-rabbit HRP conjugated secondary antibodies at 1:10,000 , incubate at RT for 1 hr.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.