

Goat IgG Texas Red™

Catalog # ASR1045

Product Information

Description	GOAT IgG whole molecule Texas Red™ conjugated
Conjugate	Texas Red®
FP Value	3.0 moles Texas Red® per mole of Goat IgG
Physical State	Lyophilized
Host Isotype	IgG
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Species of Origin	Goat
Reconstitution Volume	1.0 mL
Reconstitution Buffer	Restore with deionized water (or equivalent)
Stabilizer	10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free
Preservative	0.01% (w/v) Sodium Azide

Additional Information


Shipping Condition	Ambient
Purity	This product was prepared from normal serum by delipidation, salt fractionation, ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat IgG and anti-Goat Serum..
Storage Condition	Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Precautions Note	This product is for research use only and is not intended for therapeutic or diagnostic applications.

Background

This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms.

Images

SDS-PAGE of Goat IgG Whole Molecule Texas Red™ Conjugated (p/n ASR1045). Lane M: 5 µL Opal Prestained Marker (p/n MB-210-0500). Lane 1: Reduced Goat IgG Whole Molecule Texas Red™ (p/n ASR1045). Lane 2:



Reduced Goat IgG F(c) Fragment (p/n 005-0103). Lane 3:
 Reduced Goat IgG F(ab) Fragment (p/n 005-0105). Lane 4:
 Reduced Goat IgM Whole Molecule (p/n 005-0107). Load:
 1 μ g for IgG, F(c) and F(ab); 3 μ g for IgM.
 Predicted/Observed size: IgG at 50 and 25 kDa; F(c) at 25
 kDa; F(ab) at 25 kDa; IgM at 70 and 23 kDa. Observed F(c)
 Fragment migrates slightly higher.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.