

Human Transferrin Fluorescein

Catalog # ASR1102

Product Information

Description HUMAN TRANSFERRIN Fluorescein conjugated

Conjugate Fluorescein (FITC)

FP Value 1.68 moles Fluorescein (FITC) per mole of Human Transferrin

Application DB

Physical StateLyophilizedHost IsotypeTransferrin

Buffer 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

Species of Origin Human **Reconstitution Volume** 1.0 mL

Reconstitution Buffer Restore with deionized water (or equivalent)

Stabilizer 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free

Preservative 0.01% (w/v) Sodium Azide

Additional Information

Shipping Condition Ambient

Purity This product was prepared from normal serum by delipidation, selective

precipitation and tandem molecular sieve chromatography followed by

extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Fluorescein, anti-Human Transferrin and anti-Human Serum.

Storage Condition Store vial at 4° C prior to restoration. For extended storage aliquot

contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted

liquid. Dilute only prior to immediate use.

Precautions NoteThis product is for research use only and is not intended for therapeutic or

diagnostic applications.


Background


This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms.

Images

Dot Blot of Fluorescein conjugated Human Transferrin Antigen: Human Transferrin Fluorescein Load: 3-fold

serial dilution starting at 200 ng Block: MB-070 for 30 min at ${\rm RT}$





Dot Blot of Fluorescein conjugated Human Transferrin. Antigen: Human Transferrin Fluorescein. Load: 3-fold serial dilution starting at 200 ng. Block: None.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.