

## Anti-Rabbit IgG (H&L) Secondary Antibody

Donkey Polyclonal, Unconjugated Catalog # ASR1122

## **Product Information**

**Application** 

**Description** Anti-RABBIT IgG (H&L) (DONKEY) Antibody

**WB** 

HostDonkeyConjugateUnconjugatedTarget SpeciesRabbitClonalityPolyclonal

Physical State Liquid (sterile filtered)

Host Isotype IgG
Target Isotype IgG (H&L)

**Buffer** 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

**Immunogen** Rabbit IgG whole molecule

**Stabilizer** None

**Preservative** 0.01% (w/v) Sodium Azide

## **Additional Information**

Shipping Condition Wet Ice

**Purity** This product was prepared from monospecific antiserum by immunoaffinity

chromatography using Rabbit IgG coupled to agarose beads followed by solid

phase adsorption(s) to remove any unwanted reactivities. Assay by

immunoelectrophoresis resulted in a single precipitin arc against anti-Donkey

Serum, Rabbit IgG and Rabbit Serum.

**Storage Condition** Store vial at 4° C prior to opening. This product is stable for several weeks

at 4° C as an undiluted liquid. Dilute only prior to immediate use. For extended storage aliquot contents and freeze at -20° C or below. Avoid

cycles of freezing and thawing.

**Precautions Note** This product is for research use only and is not intended for therapeutic or

diagnostic applications.

## **Images**

Western Blot of Anti-Rabbit IgG (H&L) (DONKEY) Antibody Peroxidase Conjugated (p/n 611-7302). Lane M: 3  $\mu$ l Molecular Ladder. Lane 1: Rabbit IgG whole molecule (p/n 011-0102). Lane 2: Rabbit IgG F(ab) Fragment (p/n 011-0105). Lane 3: Rabbit IgG F(c) Fragment (p/n 010-0103). Lane 4: Rabbit IgM Whole Molecule (p/n 011-0107). Lane 5: Normal Rabbit Serum (p/n B309). All



samples were reduced. Load: 50 ng per lane. Block: MB-070 for 30 min at RT. Primary Antibody: Anti-Rabbit IgG (H&L) (DONKEY) Antibody Peroxidase Conjugated (p/n 611-7302) 1:5,000 for 60 min at RT. Secondary antibody: None. Predicted/Obsevered Size: 25 and 50 kDa for Rabbit IgG and Serum, 25 kDa for F(c) and F(ab), 70 and 23 kDa for IgM. Rabbit F(c) migrates slightly higher.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.