

Anti-Goat IgG (H&L) Secondary Antibody

Donkey Polyclonal, Unconjugated

Catalog # ASR1147

Product Information

Description	Anti-GOAT IgG (H&L) (DONKEY) Antibody
Host	Donkey
Conjugate	Unconjugated
Target Species	Goat
Reactivity	Goat
Clonality	Polyclonal
Application	WB
Physical State	Liquid (sterile filtered)
Host Isotype	IgG
Target Isotype	IgG (H&L)
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Immunogen	Anti-Goat IgG (h&L) was produced by repeated immunization with goat IgG whole molecule in donkey
Stabilizer	None
Preservative	0.01% (w/v) Sodium Azide

Additional Information


Shipping Condition	Wet Ice
Application Note	Antibody Anti-Goat IgG (H&L) is suitable for immunoblotting (western or dot blot), ELISA, immunoelectron microscopy and immunohistochemistry as well as other antibody-based enzymatic assays requiring lot-to-lot consistency.
Purity	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Goat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Donkey Serum, Goat IgG and Goat Serum.
Storage Condition	Store vial at 4° C prior to opening. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing.
Precautions Note	This product is for research use only and is not intended for therapeutic or diagnostic applications.

Background

Anti-Goat IgG (H&L) antibody generated in donkey detects specifically goat IgG. This secondary antibody anti-Goat is ideal for investigators who routinely perform titration assays, western-blot,

immunoprecipitation and more generally immunoassays.

Images



Western Blot of Donkey anti-Goat IgG antibody. Lane 1: Goat IgG. Lane 2: none. Load: 50 ng per lane. Primary antibody: Goat IgG antibody at 1:1000 for overnight at 4°C. Secondary antibody: Peroxidase donkey secondary antibody at 1:40,000 for 30 min at RT. Block: MB-070 for 30 min at RT. Predicted/Observed size: 28 and 55 kDa for Goat IgG. Other band(s): none.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.