10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

# Mouse IgG Rhodamine

Catalog # ASR1205

## Product Information

| | |
|---|---|
| **Description** | MOUSE IgG whole molecule Rhodamine conjugated |
| **Conjugate** | Rhodamine (TRITC) |
| **FP Value** | 3.5 moles Rhodamine (TRITC) per mole of Mouse IgG |
| **Physical State** | Lyophilized |
| **Host Isotype** | IgG |
| **Buffer** | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |
| **Species of Origin** | Mouse |
| **Reconstitution Volume** | 1.0 mL |
| **Reconstitution Buffer** | Restore with deionized water (or equivalent) |
| **Stabilizer** | 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free |
| **Preservative** | 0.01% (w/v) Sodium Azide |

## Additional Information

| | |
|---|---|
| **Shipping Condition** | Ambient |
| **Purity** | This product was prepared from normal serum by delipidation, salt fractionation, ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Mouse IgG and anti-Mouse Serum. |
| **Storage Condition** | Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. |
| **Precautions Note** | This product is for research use only and is not intended for therapeutic or diagnostic applications. |

## Background

This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms.

## Images

SDS-PAGE of Mouse IgG Whole Molecule Rhodamine Conjugated (p/n ASR1205). Lane 1: 5 µL Opal Prestained Marker (p/n MB-210-0500). Lane 2: Reduced Mouse IgG Whole Molecule Rhodamine Conjugated (p/n ASR1205).




Lane 3: Reduced Mouse F(c) Fragment (p/n 010-0103). Lane 4: Reduced Mouse F(ab) Fragment (p/n 010-0105). Lane 5: Mouse IgM Kappa Myeloma Protein (p/n 010-0107). Load: 1 µg per lane. Predicted/Observed size: IgG at 50 and 25 kDa; F(c) at 25 kDa; F(ab) at 25 kDa; IgM K at 70 and 23 kDa. Observed F(c) Fragment migrates slightly higher.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.