

Mouse IgG1 isotype control Biotin

Monoclonal MG1 IgG1 , Biotin

Catalog # ASR1207

Product Information

Description	MOUSE IgG1 isotype control Biotin conjugated
Conjugate	Biotin
Clonality	Monoclonal MG1 IgG1
Physical State	Lyophilized
Host Isotype	IgG1
Buffer	0.02 M Potassium Phosphate, 0.5 M Sodium Chloride, pH 7.2
Species of Origin	Mouse
Reconstitution Volume	100 μ L
Reconstitution Buffer	Restore with deionized water (or equivalent)
Stabilizer	10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free
Preservative	0.01% (w/v) Sodium Azide

Additional Information

Shipping Condition	Ambient
Application Note	Mouse IgG1 isotype control can be utilized as a control or standard reagent in Flow cytometry, Western Blotting, and ELISA experiments where determination of sample isotype is important.
Purity	This product has been prepared from in vitro cell culture by selective precipitation. In an Ouchterlony double diffusion assay the material is non-reactive with antisera to mouse IgG2a, IgG2b, IgG3, IgM, and IgA. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Mouse IgG and anti-Mouse serum. Typically, less than 1% cross reactivity was detected by ELISA against other mouse and human heavy or light chains isotypes using chain specific antibodies.
Storage Condition	Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Precautions Note	This product is for research use only and is not intended for therapeutic or diagnostic applications.

Background

Isotype control Mouse IgG1 is important for Flow Cytometry. Mouse IgG1 control has no specificity for target cells within a particular experiment. Their purpose is to confirm the specificity of primary antibody binding that it is not a result of non-specific Fc receptor binding to cells or other cellular protein interactions.

Isotype controls need to be matched to the specific primary Abs (species and isotype, including heavy and light chains) being used.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.