

## Mouse IgG3 isotype control Biotin

Monoclonal MG3 IgG3 , Biotin Catalog # ASR1210

## **Product Information**

**Description** MOUSE IgG3 isotype control Biotin conjugated

Conjugate Biotin

Clonality Monoclonal MG3 IgG3

Physical State Lyophilized Host Isotype IgG3

**Buffer** 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

Species of OriginMouseReconstitution Volume100 □

**Reconstitution Buffer** Restore with deionized water (or equivalent)

Stabilizer 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free

**Preservative** 0.01% (w/v) Sodium Azide

## **Additional Information**

Shipping Condition Ambient

**Application Note** Mouse IgG3 isotype control can be utilized as a control or standard reagent

in Flow cytometry, Western Blotting, and ELISA experiments where

determination of sample isotype is important.

**Purity** This product has been prepared from in vitro cell culture by selective

precipitation. In an Ouchterlony double diffusion assay the material is non-reactive with antisera to mouse IgG1, IgG2a, IgG2b, IgM, and IgA. Assay

by immunoelectrophoresis resulted in a single precipitin arc against

anti-Mouse IgG and anti-Mouse serum. Typically, less than 1% cross reactivity was detected by ELISA against other mouse isotypes using chain specific

antibodies.

**Storage Condition** Store vial at 4° C prior to restoration. For extended storage aliquot

contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted

liquid. Dilute only prior to immediate use.

**Precautions Note**This product is for research use only and is not intended for therapeutic or

diagnostic applications.

## **Background**

Isotype controls are important for Flow Cytometry and have no specificity for target cells within a particular experiment. Their purpose is to confirm the specificity of primary antibody binding that it is not a result of non-specific Fc receptor binding to cells or other cellular protein interactions. Isotype controls need to be

matched to the specific primary Abs (species and isotype, including heavy and light chains) being used.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.