

Anti-Human IgG F(c) Secondary Antibody

Rabbit Polyclonal, Unconjugated

Catalog # ASR1348

Product Information


Description	Anti-HUMAN IgG F(c) (RABBIT) Antibody
Host	Rabbit
Conjugate	Unconjugated
Target Species	Human
Clonality	Polyclonal
Application	WB
Physical State	Lyophilized
Host Isotype	Antiserum
Target Isotype	IgG F(c)
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Immunogen	Human IgG F(c) fragment
Reconstitution Volume	2.0 mL
Reconstitution Buffer	Restore with deionized water (or equivalent)
Stabilizer	None
Preservative	0.01% (w/v) Sodium Azide

Additional Information

Shipping Condition	Ambient
Purity	Human IgG F(c) Antibody was prepared from monospecific antiserum by a delipidation and defibrination. Assay by immunoelectrophoresis resulted in a single precipitin arc against Human IgG, Human IgG F(c) and Human Serum. No reaction was observed against Human IgG F(ab') ₂ .
Storage Condition	Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Precautions Note	This product is for research use only and is not intended for therapeutic or diagnostic applications.

Images

Western Blot of Rabbit anti-Human IgG F(c) Secondary Antibody. Lane 1: Human F(c). Lane 2: None. Load: 50 ng per lane. Primary antibody: Human IgG F(c) secondary antibody at 1:1000 overnight at 4°C. Secondary antibody: DyLight™ 649 rabbit secondary antibody at 1:20,000 for



30 min at RT. Block: MB-070 for 30 min at RT.
Predicted/Observed size: 28 kDa, 28 kDa for Human IgG
F(c). Other band(s): None.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.