

## Chicken IgG Peroxidase

Catalog # ASR1455

## **Product Information**

**Description** CHICKEN IgG whole molecule Peroxidase conjugated

**Conjugate** Peroxidase (Horseradish)

Physical State Lyophilized

Host Isotype IgG

**Buffer** 0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2

**Species of Origin** Chicken **Reconstitution Volume** 1.0 mL

**Reconstitution Buffer** Restore with deionized water (or equivalent)

Stabilizer 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free

**Preservative** None


## **Additional Information**

<b>Shipping Condition</b>	Ambient
Purity	This product was prepared from normal serum by delipidation, salt fractionation, ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti- Chicken IgG and anti-Chicken Serum.
Storage Condition	Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

**Precautions Note** This product is for research use only and is not intended for therapeutic or

diagnostic applications.

## **Images**



SDS-PAGE of Chicken IgG Whole Molecule Peroxidase Conjugated (p/n ASR1455). Lane M: 5 µL Opal Prestained Marker (p/n MB-210-0500). Lane 1: Reduced Chicken IgG Whole Molecule Peroxidase Conjugated (p/n ASR1455). Lane 2: Reduced Chicken IgG F(c) Fragment (p/n 003-0103). Lane 3: Reduced Chicken IgG F(ab) Fragment (p/n 003-0105). Lane 4: Reduced Chicken IgM Whole Molecule (p/n 003-0107). Load: 1 µg per lane.

Predicted/Observed size: IgG at 72 and 25 kDa; F(c) at 25 kDa; F(ab) at 25 kDa; IgM at 75 kDa. Observed F(c)

Fragment migrates slightly higher. Other bands: Chicken IgG heavy chain alternative splicing variant at approximately 40 kDa in Lane 1.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.