

Mouse IgG Biotin

Catalog # ASR1495

Product Information

Description MOUSE IgG whole molecule Biotin conjugated

ConjugateBiotinPhysical StateLyophilized

Host Isotype IgG

Buffer 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

Species of Origin Mouse **Reconstitution Volume** 1.0 mL

Reconstitution Buffer Restore with deionized water (or equivalent)

Additional Information

Shipping Condition	Ambient
--------------------	---------

Purity This product was prepared from normal serum delipidation, salt

fractionation, ion exchange chromatography followed by papain digestion

and extensive dialysis against the buffer stated above. Assay by

immunoelectrophoresis resulted in a single precipitin arc against anti-biotin,

anti-Mouse IgG and anti-Mouse Serum.

Storage Condition Store vial at 4° C prior to restoration. For extended storage aliquot


contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted

liquid. Dilute only prior to immediate use.

Precautions NoteThis product is for research use only and is not intended for therapeutic or

diagnostic applications.

Images



SDS-PAGE of Mouse IgG Whole Molecule Biotin Conjugated (p/n ASR1495). Lane 1: 5 μ L Opal Prestained Marker (p/n MB-210-0500). Lane 2: Reduced Mouse IgG Whole Molecule Biotin Conjugated (p/n ASR1495). Lane 3: Reduced Mouse F(c) Fragment (p/n 010-0103). Lane 4: Reduced Mouse F(ab) Fragment (p/n 010-0105). Lane 5: Mouse IgM Kappa Myeloma Protein (p/n 010-0107). Load: 1 μ g per lane. Predicted/Observed size: IgG at 50 and 25 kDa; F(c) at 25 kDa; F(ab) at 25 kDa; IgM K at 70 and 23 kDa. Observed F(c) Fragment migrates slightly higher.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.