10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

# Rabbit IgG Fab Biotin

Catalog # ASR1501


## Product Information

| | |
|---|---|
| **Description** | RABBIT IgG F(ab) fragment Biotin conjugated |
| **Conjugate** | Biotin |
| **Physical State** | Lyophilized |
| **Host Isotype** | IgG F(ab) |
| **Buffer** | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |
| **Species of Origin** | Rabbit |
| **Reconstitution Volume** | 1.0 mL |
| **Reconstitution Buffer** | Restore with deionized water (or equivalent) |
| **Stabilizer** | 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free |
| **Preservative** | 0.01% (w/v) Sodium Azide |

## Additional Information

| | |
|---|---|
| **Shipping Condition** | Ambient |
| **Purity** | This product was prepared from normal serum by delipidation, salt fractionation, ion exchange chromatography followed by papain digestion and extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-biotin, anti-Rabbit IgG, anti-Rabbit IgG F(ab')2 and anti-Rabbit Serum. No reaction was observed against anti-Rabbit IgG F(c) or anti- Papain. |
| **Storage Condition** | Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. |
| **Precautions Note** | This product is for research use only and is not intended for therapeutic or diagnostic applications. |

## Images




SDS-PAGE of Rabbit IgG F(ab) Fragment Biotin Conjugated (p/n ASR1501). Lane M: 3 µL Opal Prestained Marker (p/n MB-210-0500). Lane 1: Reduced Rabbit IgG Whole Molecule (p/n 011-0102). Lane 2: Reduced Rabbit IgG F(ab) Fragment Biotin Conjugated (p/n ASR1501). Lane 3: Reduced Rabbit IgG F(c) Fragment (p/n 011-0103). Lane 4: Reduced Rabbit IgM Whole Molecule (p/n 011-0107). Load: 1 µg for F(ab) and F(c); 1.2 µg for IgG and IgM. Predicted/Observed size: IgG at 50 and 25 kDa; F(c) at 25

kDa; F(ab) at 25 kDa; IgM at 70 and 23 kDa. Observed F(c) Fragment migrates slightly higher.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.