10320 Camino Santa Fe, Suite G San Diego, CA 92121 Tel: 858.875.1900 Fax: 858.875.1999



Anti-Human IgG (H&L) (Biotin Conjugated) Secondary Antibody

Donkey Polyclonal, Biotin Catalog # ASR1610

Product Information

Description Anti-HUMAN IgG (H&L) (DONKEY) Antibody Biotin Conjugated

HostDonkeyConjugateBiotinTarget SpeciesHumanClonalityPolyclonalPhysical StateLyophilized

Host Isotype IgG

Target Isotype IgG (H&L)

Buffer 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

Immunogen Human IgG whole molecule

Reconstitution Volume 1.0 mL

Reconstitution Buffer Restore with deionized water (or equivalent)

Stabilizer 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free

Preservative 0.01% (w/v) Sodium Azide

Additional Information

Shipping Condition Ambient

Application Note This product has been assayed against 1.0 ug of Human IgG in a standard

capture ELISA using Peroxidase Conjugated Streptavidin #S000-03 and ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of

1:30,000 to 1:130,000 is suggested for this product.

Purity This product was prepared from monospecific antiserum by immunoaffinity

chromatography using Human IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-biotin,

anti-Donkey Serum, Human IgG and Human Serum.

Storage Condition Store vial at 4° C prior to restoration. For extended storage aliquot

contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted

liquid. Dilute only prior to immediate use.

Precautions Note This product is for research use only and is not intended for therapeutic or

diagnostic applications.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.