10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

# Anti-Rabbit IgG (H&L) (Texas Red™ Conjugated) Secondary Antibody

Goat Polyclonal, Texas Red®
Catalog # ASR1635

## Product Information

| | |
|---|---|
| **Description** | Anti-RABBIT IgG (H&L) (GOAT) Antibody Texas Red™ Conjugated |
| **Host** | Goat |
| **Conjugate** | Texas Red® |
| **FP Value** | 1.9 moles Texas Red® per mole of IgG |
| **Target Species** | Rabbit |
| **Clonality** | Polyclonal |
| **Application** | WB, DB |
| **Physical State** | Lyophilized |
| **Host Isotype** | IgG |
| **Target Isotype** | IgG (H&L) |
| **Buffer** | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |
| **Immunogen** | Rabbit IgG whole molecule |
| **Reconstitution Volume** | 1.0 mL |
| **Reconstitution Buffer** | Restore with deionized water (or equivalent) |
| **Stabilizer** | 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free |
| **Preservative** | 0.01% (w/v) Sodium Azide |


## Additional Information

| | |
|---|---|
| **Shipping Condition** | Ambient |
| **Purity** | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Rabbit IgG coupled to agarose. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Rabbit IgG and Rabbit Serum. |
| **Storage Condition** | Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. |
| **Precautions Note** | This product is for research use only and is not intended for therapeutic or diagnostic applications. |


## Background

This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms.


## Images




Western Blot of Anti-Rabbit IgG (H&L) (GOAT) Antibody (p/n 611-1102). Lane M: 3 µl Molecular Ladder. Lane 1: Rabbit IgG whole molecule (p/n 011-0102). Lane 2: Rabbit IgG F(ab) Fragment (p/n 011-0105). Lane 3: Rabbit IgG F(c) Fragment (p/n 010-0103). Lane 4: Rabbit IgM Whole Molecule (p/n 011-0107). Lane 5: Normal Rabbit Serum (p/n B309). All samples were reduced. Load: 50 ng per lane. Block: MB-070 for 30 min at RT. Primary Antibody: Anti-Rabbit IgG (H&L) (GOAT) Antibody (p/n 611-1102) 1:1,000 for 60 min at RT. Secondary antibody: Anti-Goat IgG (DONKEY) Peroxidase Conjugated Antibody (p/n CUST10) 1:40,000 in MB-070 for 30 min at RT. Predicted/Obsevered Size: 25 and 50 kDa for Rabbit IgG and Serum, 25 kDa for F(c) and F(ab), 70 and 23 kDa for IgM. Rabbit F(c) migrates slightly higher.




Dot Blot showing the detection of Rabbit IgG. A three-fold serial dilution of Rabbit IgG starting at 200ng was spotted onto 0.45 ?m nitrocellulose. After blocking in 5% Blotto (p/n B501-0500) 1 Hour at 20°C, Anti-Rabbit IgG (H&L) (GOAT) Antibody Texas Red Conjugated (p/n ASR1635) secondary antibody was used at 1:1000 in Blocking Buffer for Fluorescent Western Blotting (p/n MB-070) and imaged using the Bio-Rad VersaDoc® 4000 MP.




Western blot of Texas Red™ Conjugated Goat Anti-Rabbit IgG secondary antibody. Lane 1: Rabbit IgG. Lane 2: None. Load: 50 ng per lane. Primary antibody: None. Secondary antibody: Texas Red™ goat secondary antibody at 1:1,000 for 60 min at RT. Blocking: MB-070 for 30 min at RT. Predicted/Observed size: 25 & 55 kDa, 25 & 55 kDa for Rabbit IgG. Other band(s): None.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.