10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

# F(ab')2 Anti-Guinea Pig IgG (H&L) Secondary Antibody

Rabbit Polyclonal, Unconjugated
Catalog # ASR1688

## Product Information

| | |
|---|---|
| **Description** | F(ab')2 Anti-GUINEA PIG IgG [H&L] (RABBIT) Antibody |
| **Host** | Rabbit |
| **Conjugate** | Unconjugated |
| **Target Species** | Guinea Pig |
| **Clonality** | Polyclonal |
| **Physical State** | Liquid (sterile filtered) |
| **Host Isotype** | IgG F(ab')2 |
| **Target Isotype** | IgG (H&L) |
| **Buffer** | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |
| **Immunogen** | Guinea Pig IgG whole molecule |
| **Stabilizer** | None |

## Additional Information

| | |
|---|---|
| **Shipping Condition** | Wet Ice |
| **Application Note** | Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. The maximum amount of reagent required to stain 1 x 10E6 cells in flow cytometry is approximately 1.0 □g of antibody. Lesser amounts of reagent may be sufficient for staining. Optimal titers for other applications should be determined by the researcher. As a general guideline dilutions of 1:100 to 1:250 should be suitable for most applications. |
| **Purity** | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Guinea Pig IgG coupled to agarose beads followed by pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum, Guinea Pig IgG and Guinea Pig Serum. No reaction was observed against anti-Pepsin and anti-Rabbit IgG F(c). |
| **Storage Condition** | Store vial at 4° C prior to opening. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. |
| **Precautions Note** | This product is for research use only and is not intended for therapeutic or diagnostic applications. |

## Background

F(ab')2 Antibody was generated by enzymatic cleavage and subsequent separation from the Fc fragment. Because of their smaller size, F(ab)2 fragments offer several advantages over intact antibodies for use in certain immunochemical techniques and experimental applications. F(ab)2 fragments penetrate into tissue samples and show better antigen recognition and signal generation in IHC. F(ab)2 fragments lack the Fc region and therefore do not bind Fc receptors which effectively lowers background staining. F(ab')2 Antibody is ideal for investigators who routinely perform flow cytometry, immunohistochemistry or IHC and other immunoassays.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.