10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

F(ab')2 Anti-Human IgG IgA IgM (H&L) Pre-Adsorbed Secondary Antibody

Goat Polyclonal, Unconjugated
Catalog # ASR1735

Product Information

Description	F(ab')2 Anti-HUMAN IgG IgA IgM (H&L) (GOAT) Antibody Min X MOUSE Serum Proteins
Host	Goat
Conjugate	Unconjugated
Target Species	Human
Clonality	Polyclonal
Physical State	Liquid (sterile filtered)
Host Isotype	IgG F(ab')2
Target Isotype	IgG IgA IgM
Buffer	0.125 M Sodium Borate, 0.075 M Sodium Chloride, 0.005 M EDTA, pH 8.0
Immunogen	Human IgG, IgA, and IgM whole molecules
Stabilizer	None
Preservative	0.01% (w/v) Sodium Azide

Additional Information

Shipping Condition	Wet Ice
Application Note	Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. The maximum amount of reagent required to stain 1 x 10E6 cells in flow cytometry is approximately 1.0 µg of antibody. Lesser amounts of reagent may be sufficient for staining. Optimal titers for other applications should be determined by the researcher. As a general guideline dilutions of 1:100 to 1:250 should be suitable for most applications.
Purity	This product was prepared from polyspecific antiserum by immunoaffinity chromatography using antigens coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum. No reaction was observed against anti-Goat IgG F(c), anti-Pepsin or Mouse Serum Proteins. This product is suitable for the detection of all Human immunoglobulin classes, isotypes and chain combinations.
Storage Condition	Store vial at 4° C prior to opening. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing.
Precautions Note	This product is for research use only and is not intended for therapeutic or

diagnostic applications.

Background

F(ab')2 Antibody was generated by enzymatic cleavage and subsequent separation from the Fc fragment. Because of their smaller size, F(ab)2 fragments offer several advantages over intact antibodies for use in certain immunochemical techniques and experimental applications. F(ab)2 fragments penetrate into tissue samples and show better antigen recognition and signal generation in IHC. F(ab)2 fragments lack the Fc region and therefore do not bind Fc receptors which effectively lowers background staining. F(ab')2 Antibody is ideal for investigators who routinely perform flow cytometry, immunohistochemistry or IHC and other immunoassays.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.