

Anti-Bovine IgG F(c) Secondary Antibody

Rabbit Polyclonal, Unconjugated Catalog # ASR1751

Product Information

Description Anti-BOVINE IgG F(c) (RABBIT) Antibody

Host Rabbit

Conjugate Unconjugated

Target Species Bovine Reactivity Bovine Clonality Polyclonal

Physical State Liquid (sterile filtered)

Host Isotype IgG **Target Isotype** IgG F(c)

Buffer 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

Anti-Bovine IgG F(c) was produced by repeated immunization with bovine **Immunogen**

IgG F(c) fragment in rabbit.

Stabilizer None

Preservative 0.01% (w/v) Sodium Azide

Additional Information

Shipping Condition Wet Ice

Application Note Bovine IgG F(c) Antibody is suitable for immunoblotting (western or dot

blot), ELISA, and immunohistochemistry requiring extremely low background

levels, lot-to-lot consistency, high titer and specificity.

This product was prepared from monospecific antiserum by immunoaffinity **Purity**

chromatography using Bovine IgG coupled to agarose beads followed by solid

phase adsorption(s) to remove any unwanted reactivities. Assay by

immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum, Bovine IgG, Bovine IgG F(c) and Bovine Serum. No reaction was

observed against Bovine IgG F(ab')2.

Store vial at 4° C prior to opening. This product is stable for several weeks **Storage Condition**

at 4° C as an undiluted liquid. Dilute only prior to immediate use. For

extended storage aliquot contents and freeze at -20° C or below. Avoid cycles

of freezing and thawing.

Precautions Note This product is for research use only and is not intended for therapeutic or

diagnostic applications.

Background

Anti-Bovine IgG F(c) antibody generated in rabbit detects specifically bovine IgG. This secondary antibody anti-Bovine is ideal for investigators who routinely perform titration assays, western-blot,

 $immun oprecipitation\ and\ more\ generally\ immunoassays.$

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.