

Anti-Human IgM (Biotin Conjugated) Secondary Antibody

Goat Polyclonal, Biotin
Catalog # ASR1780

Product Information

Description	Anti-HUMAN IgM (GOAT) Antibody Biotin Conjugated
Host	Goat
Conjugate	Biotin
Target Species	Human
Clonality	Polyclonal
Physical State	Lyophilized
Host Isotype	IgG
Target Isotype	IgM μ chain
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Immunogen	Human IgM mu heavy chain
Species of Origin	Human
Reconstitution Volume	1.0 mL
Reconstitution Buffer	Restore with deionized water (or equivalent)
Stabilizer	10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free
Preservative	0.01% (w/v) Sodium Azide

Additional Information

Shipping Condition Ambient

Application Note Anti-HUMAN IgM (μ chain) (GOAT) Antibody has been assayed against 1.0 μ g of Human IgM in a standard capture ELISA using Peroxidase Conjugated Streptavidin #S000-03 and ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:3,000 to 1:17,000 of the reconstitution concentration is suggested for this product. This product has been assayed against 1.0 μ g of Human IgM in a standard capture ELISA using Peroxidase Conjugated Streptavidin #S000-03 and ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:3,000 to 1:17,000 of the reconstitution concentration is suggested for this product. Anti-HUMAN IgM (μ chain) (GOAT) Antibody has been assayed against 1.0 μ g of Human IgM in a standard capture ELISA using Peroxidase Conjugated Streptavidin #S000-03 and ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:3,000 to 1:17,000 of the reconstitution concentration is suggested for this product.

Purity Anti-HUMAN IgM (μ chain) (GOAT) Antibody was prepared from monospecific antiserum by immunoaffinity chromatography using Human IgM coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-biotin, anti-Goat Serum, Human

IgM and Human Serum. No reaction was observed against other Human heavy or light chain proteins.

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Background

Human IgM alpha heavy chain antibody generated in goat detects specifically Human IgM alpha heavy chain. Anti-Human IgM antibody is ideal for investigators who routinely perform ELISA, Sandwich ELISA, titration assays, western-blot, immunoprecipitation and more generally immunoassays.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.