10320 Camino Santa Fe, Suite G San Diego, CA 92121 Tel: 858.875.1900 Fax: 858.875.1999



F(ab')2 Anti-Goat IgG (H&L) (Fluorescein Conjugated) Pre-Adsorbed Secondary Antibody

Donkey Polyclonal, Fluorescein (FITC) Catalog # ASR2079

Product Information

Description F(ab')2 Anti-GOAT IgG [H&L] (DONKEY) Antibody Fluorescein

Conjugated Min X Ch GP Ham Hs Hu Ms Rb & Rt Serum Proteins

Host Donkey

Conjugate Fluorescein (FITC)

FP Value 5. moles Fluorescein (FITC) per mole of IgG F(ab')2

Target SpeciesGoatClonalityPolyclonalPhysical StateLyophilizedHost IsotypeIgG F(ab')2

Target Isotype IgG (H&L)

Buffer 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

Immunogen Goat IgG whole molecule

Reconstitution Volume 500 □

Reconstitution Buffer Restore with deionized water (or equivalent)

Stabilizer 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free

Preservative 0.01% (w/v) Sodium Azide

Additional Information

Shipping Condition Ambient

Application Note Suitable for immunomicroscopy and flow cytometry or FACS analysis as well

as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency,

high titer and specificity.

Purity This product was prepared from monospecific antiserum by immunoaffinity

chromatography using Goat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis

resulted in a single precipitin arc against anti-Fluorescein, anti-Donkey Serum, Goat IgG and Goat Serum. No reaction was observed against anti-Pepsin, anti-Donkey IgG F(c) or Chicken, Guinea Pig, Hamster, Horse, Human, Mouse,

Rabbit and Rat Serum Proteins.

Storage Condition Store vial at 4° C prior to restoration. For extended storage aliquot

contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted

liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Background

This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.