

Anti-Ferret IgA (Biotin Conjugated) Secondary Antibody

Goat Polyclonal, Biotin Catalog # ASR2121

Product Information

Description Anti-FERRET IgA(GOAT) Antibody Biotin Conjugated

HostGoatConjugateBiotinTarget SpeciesFerretClonalityPolyclonalPhysical StateLyophilized

Host Isotype IgG
Target Isotype IgA

Buffer 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

Immunogen Ferret IgA alpha heavy chain

Species of Origin Ferret **Reconstitution Volume** 1.0 mL

Reconstitution Buffer Restore with deionized water (or equivalent)

Stabilizer 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free

Preservative 0.01% (w/v) Sodium Azide

Additional Information

Shipping Condition Ambient

Application Note Anti-FERRET IgA (alpha chain) (GOAT) Antibody has been assayed against 1.0

ug of Ferret IgA in a standard capture ELISA using Peroxidase Conjugated

Streptavidin #S000-03 and ABTS

(2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:5,000

to 1:20,000 is suggested for this product.

Purity Anti-FERRET IgA (alpha chain) (GOAT) Antibody was prepared from

monospecific antiserum by immunoaffinity chromatography using Ferret IgA coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-biotin, anti-Goat Serum, Ferret IgA and Ferret

Serum. No reaction was observed against Ferret IgM or Ferret IgG.

Storage Condition Store vial at 4° C prior to restoration. For extended storage aliquot

contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted

liquid. Dilute only prior to immediate use.

Precautions NoteThis product is for research use only and is not intended for therapeutic or

diagnostic applications.

Background

Ferret IgA whole molecule antibody generated in goat detects specifically ferret IgA whole molecule. Anti-Human IgA is ideal for investigators who routinely perform ELISA, Sandwich ELISA, titration assays, western-blot, immunoprecipitation and more generally immunoassays

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.